NATIONAL CANCER CONTROL PROGRAMME (NCCP) 00720 · VERSION 2

Neratinib · extended adjuvant treatment

Neratinib is taken once daily with food for one year. The usual starting dose is 240 mg; the team may interrupt or reduce it for toxicity. Treatment starts less than one year after completing adjuvant trastuzumab.

Your treatment and review passport

Bring this to reviews. Record symptoms, treatment changes and questions when helpful. You do not need to complete a daily diary or calculate a toxicity grade.

The four sheets are reusable review periods, not a prescribed number of cycles. Follow the treatment and rest days agreed with your team; do not calculate a restart from this sheet.

These patient-language prompts are not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) assessment. Urgent symptoms override the form.

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Report diarrhoea early. Poor intake, dizziness, reduced urine, blood in stool, fever, severe pain or jaundice needs prompt clinical advice.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

Your complete treatment reference

https://sactcheck.com/patient/00720/

Public information only. No personal information is encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Neratinib · extended adjuvant treatment

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	☐	☐	☐	
Nausea / vomiting	☐	☐	☐	
Hydration / urine	☐	☐	☐	
Fatigue	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Report diarrhoea early. Poor intake, dizziness, reduced urine, blood in stool, fever, severe pain or jaundice needs prompt clinical advice. Emergency symptoms: 112 / 999.

Neratinib · extended adjuvant treatment

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	□	□	□	
Nausea / vomiting	□	□	□	
Hydration / urine	□	□	□	
Fatigue	□	□	□	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Report diarrhoea early. Poor intake, dizziness, reduced urine, blood in stool, fever, severe pain or jaundice needs prompt clinical advice. Emergency symptoms: 112 / 999.

Neratinib · extended adjuvant treatment

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	□	□	□	
Nausea / vomiting	□	□	□	
Hydration / urine	□	□	□	
Fatigue	□	□	□	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Report diarrhoea early. Poor intake, dizziness, reduced urine, blood in stool, fever, severe pain or jaundice needs prompt clinical advice. Emergency symptoms: 112 / 999.

Neratinib · extended adjuvant treatment

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	□	□	□	
Nausea / vomiting	□	□	□	
Hydration / urine	□	□	□	
Fatigue	□	□	□	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Report diarrhoea early. Poor intake, dizziness, reduced urine, blood in stool, fever, severe pain or jaundice needs prompt clinical advice. Emergency symptoms: 112 / 999.