# TCHP: docetaxel, carboplatin, trastuzumab and pertuzumab

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Day 1 · carboplatin combination

**Active medicines:** Docetaxel + Carboplatin + Trastuzumab + Pertuzumab

**Emetogenic risk:** High · carboplatin AUC ≥4

| Medicine | Dose / route | When |
|---|---|---|
| Aprepitant | 125 mg orally | Day 1 |
| Ondansetron | 16 mg orally | Day 1<br>Individual dose / QT / hepatic review |
| Dexamethasone | 12 mg orally | Day 1<br>Reconcile chemotherapy premedication steroids |
| Aprepitant | 80 mg orally daily | Days 2–3 |
| Metoclopramide | 10 mg orally up to three times daily | Subsequent days: duration and use to be confirmed<br>Individual contraindications, renal function and duration review |
| Dexamethasone | 8 mg orally daily | Days 2–4<br>Carboplatin AUC ≥4 antiemetic row; reconcile total steroid exposure |

### Additional phase-specific support

- Docetaxel: dexamethasone 8 mg orally twice daily for 3 days starting the day before treatment; minimum 3 doses pre-treatment unless contraindicated. If oral doses were missed, any 20 mg intravenous (IV) replacement is a consultant decision.
- Reconcile docetaxel premedication with the antiemetic steroid course and interactions; issue one coherent steroid prescription, not duplicate courses.
- Consider primary granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) using the source and individual risk; product and timing follow the prescribed plan.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00722 v2 · p7 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-21 · SACTCheck v0.84.16 · Regimen-level drafts; prescriber and oncology-pharmacy verification required