Pertuzumab + trastuzumab maintenance

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Pertuzumab + trastuzumab maintenance

Active medicines: Pertuzumab + Trastuzumab

Emetogenic risk: Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine prophylactic antiemetic, corticosteroid or granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) order is added for these antibodies alone. Review individual symptoms and previous infusion reactions.
- Confirm the last administration date. A gap of 6 weeks or longer requires review of both antibody loading doses under the National Cancer Control Programme (NCCP) schedule.
- Keep the existing cardiac-monitoring clock when moving to maintenance; review new breathlessness, swelling, fatigue or rapid weight gain.
- Report diarrhoea early; agree hydration, symptom treatment and when to contact the unit.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00726 v3a · p4 · NCCP antiemetics v8, Table 1 pp4–5