BASELINE → TREATMENT → FOLLOW-UP

Tests, symptoms and treatment decisions

Medicine-safety checks, tumour-tissue testing and response imaging answer different questions. Product-information additions are labelled. No universal blood tumour-marker panel or scan interval is implied.

Check	Before starting	During treatment	Between visits	Next treatment / response	After a pause or stop
Bloods / medicine safety	Blood count, kidney/liver tests, glucose, urine protein, blood pressure and cardiac history/ examination; National Cancer Control Programme (NCCP) lung testing for patients at high risk. Current lomustine SmPC additionally advises baseline lung-function assessment.	Regular blood count, kidney/liver tests and urine protein; blood pressure before each bevacizumab dose and after treatment. Current lomustine SmPC advises weekly blood counts for at least six weeks after a dose and continuing lung assessment.	Keep additional tests even on days without an anticancer dose.	Team reviews current and previous results, symptoms and dose history.	Agree continuing checks; lomustine effects may persist.
Neurology / medicines / function	Record usual symptoms, medicines, daily function and support needs.	Review headache, seizures, speech, strength, cognition and any steroid plan.	Report new changes early; do not wait for an magnetic resonance imaging (MRI).	Discuss treatment tolerance, priorities and the next authorised dose.	Agree rehabilitation, symptom support and contacts.
Response / appointments	Confirm exact diagnosis and baseline imaging.	Imaging and clinical review follow the specialist plan.	Record bookings, result-review contacts and questions.	Interpret scans with symptoms, treatment effects and medicines.	Agree the next imaging and clinical review.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00742 v3 · current regimen · Lomustine medac current product information · January 2026 · EMA Avastin product information and patient leaflet

THE WHOLE TREATMENT JOURNEY

Lomustine + bevacizumab 5 mg/kg: treatment and checks

NCCP 00742: recurrent malignant glioblastoma; bevacizumab use is off-label.

Each 42-day cycle has lomustine capsules taken once on Day 1 and bevacizumab infusions on Days 1, 15 and 29. Up to six cycles are planned, with review before each treatment.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Next treatment / response	After a pause or stop
Bloods / medicine safety NCCP 00742	Blood count, kidney/liver tests, glucose, urine protein, blood pressure and cardiac history/ examination; NCCP lung testing for patients at high risk. Current lomustine SmPC additionally advises baseline lung-function assessment.	Regular blood count, kidney/liver tests and urine protein; blood pressure before each bevacizumab dose and after treatment. Current lomustine SmPC advises weekly blood counts for at least six weeks after a dose and continuing lung assessment.	Keep additional tests even on days without an anticancer dose.	Team reviews current and previous results, symptoms and dose history.	Agree continuing checks; lomustine effects may persist.
Neurology / medicines / function NCCP 00742	Record usual symptoms, medicines, daily function and support needs.	Review headache, seizures, speech, strength, cognition and any steroid plan.	Report new changes early; do not wait for an MRI.	Discuss treatment tolerance, priorities and the next authorised dose.	Agree rehabilitation, symptom support and contacts.
Response / appointments NCCP 00742	Confirm exact diagnosis and baseline imaging.	Imaging and clinical review follow the specialist plan.	Record bookings, result-review contacts and questions.	Interpret scans with symptoms, treatment effects and medicines.	Agree the next imaging and clinical review.

What each check is for

1 Lomustine + bevacizumab 5 mg/kg

42-day cycle

Lomustine chemotherapy and bevacizumab targeted therapy are used together to help control recurrent glioblastoma. Discuss the expected benefit, symptom goals and treatment burden with your team.

Ask / act: Confirm the current phase, prescription, blood tests and support medicines.

Sources: NCCP 00742

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-05 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.39