SACTCHECK · TREATMENT PASSPORT

# Dose-dense AC → weekly paclitaxel + 3-weekly trastuzumab

## A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Four AC cycles 14 days apart, followed by paclitaxel weekly for 12 doses and intravenous (IV) trastuzumab every 21 days. Trastuzumab then continues to complete one year total.

The four pages are reusable recording sheets, not a prescribed four-cycle course. Ask for additional sheets as needed.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00745/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

# Dose-dense AC → weekly paclitaxel + 3-weekly trastuzumab

Cycle number: _____ Start date: _____ Next review: _____

**Phase 1** · **AC** · **reusable record** · copy this sheet for another cycle of the same phase.

### Doxorubicin · Anthracycline chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Mouth soreness / ulcers** | □ None | □ Mild soreness; eating normally | □ Painful; soft food needed: call | □ Cannot eat / drink enough: urgent review | □ Major breathing difficulty: 112/999 |
| **Nausea** | □ None | □ Queasy but eating and drinking | □ Intake reduced: contact the team | □ Unable to maintain intake: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

### Cyclophosphamide · Alkylating chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Nausea** | □ None | □ Queasy but eating and drinking | □ Intake reduced: contact the team | □ Unable to maintain intake: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |
| **Urinary symptoms** | □ Usual pattern | □ Mild change: report | □ Pain or frequent urination: call | □ Cannot pass urine / marked bleeding: urgent review | □ Collapse or severe illness: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Dose-dense AC → weekly paclitaxel + 3-weekly trastuzumab

Cycle number: _____ Start date: _____ Next review: _____

**Phase 2 · Paclitaxel with HER2 treatment · reusable record** · copy this sheet for another cycle of the same phase.

### Paclitaxel · Taxane chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Numbness / tingling** | □ None | □ Mild change; usual tasks unaffected | □ Limits buttons, writing or walking: call | □ Washing, dressing or walking difficult: urgent review | □ Sudden major weakness / paralysis: emergency help |
| **Muscle / joint pain** | □ None | □ Mild aches; normal activity | □ Pain limits tasks / sleep: call | □ Pain limits self-care: urgent review | □ Sudden severe deterioration: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |
| **Mouth soreness / ulcers** | □ None | □ Mild soreness; eating normally | □ Painful; soft food needed: call | □ Cannot eat / drink enough: urgent review | □ Major breathing difficulty: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Nausea** | □ None | □ Queasy but eating and drinking | □ Intake reduced: contact the team | □ Unable to maintain intake: urgent review | □ Collapse / severe dehydration: 112/999 |

### Trastuzumab · HER2-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Dose-dense AC → weekly paclitaxel + 3-weekly trastuzumab

Cycle number: _____ Start date: _____ Next review: _____

**Review** / **HER2 continuation** · **reusable record** · copy this sheet for another cycle of the same phase.

### Trastuzumab · HER2-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Dose-dense AC → weekly paclitaxel + 3-weekly trastuzumab

Cycle number: _____ Start date: _____ Next review: _____

**Review** / **HER2 continuation** · **reusable record** · copy this sheet for another cycle of the same phase.

### Trastuzumab · HER2-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.