Phesgo maintenance

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Phesgo-only maintenance

Active medicines: Pertuzumab + trastuzumab (Phesgo)

Emetogenic risk: Minimal for Phesgo / trastuzumab alone

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine prophylactic antiemetic or cytotoxic premedication is carried over. Previous reactions or symptoms need an individual plan.
- Phesgo premedication is not usually required; consider paracetamol/antihistamine for the individual reaction history.
- After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.
- Continue cardiac monitoring; maintenance does not restart the echo clock.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00785 v3 · p4 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-21 · SACTCheck v0.84.15 · Regimen-level drafts; prescriber and oncology-pharmacy verification required