# Bevacizumab + fluorouracil / folinic acid

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Bevacizumab + fluorouracil/folinic acid cycle

**Active medicines:** Bevacizumab + Folinic acid + Fluorouracil

**Emetogenic risk:** Low (fluorouracil); bevacizumab minimal

| Medicine | Dose / route | When |
|---|---|---|
| Ondansetron | 8 mg orally once | Before chemotherapy if prescribed<br>NCIS V8 low-risk default; review QT risk, interactions and constipation. Regimen premedication is not otherwise routinely required. |
| Metoclopramide | 10 mg orally up to three times daily as required | Rescue only; prescriber confirms duration<br>Review renal function, neurological contraindications and interactions. |

### Additional phase-specific support

- This regimen has a low risk of sickness, but your own symptoms still matter. Follow the prescribed prevention and rescue medicines. Call if vomiting prevents drinking; do not duplicate steroids or anti-sickness doses.
- Growth-factor injections depend on your risk, blood counts and previous complications. If prescribed, confirm their timing in relation to chemotherapy and pump removal; do not assume everyone receives them.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00791 v2 · NCCP NCIS antiemetics V8 low-risk row

Sources checked 2026-11-04 · SACTCheck v0.84.33 · Regimen-level drafts; prescriber and oncology-pharmacy verification required