# Nivolumab, pemetrexed and cisplatin before surgery

National Cancer Control Programme (NCCP) 00849 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

## Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

### Before the visit

Confirm blood tests, transport, medicine instructions and the oncology contact number.

STEP 2

### Time in the unit

Nivolumab is infused over 30 minutes before chemotherapy. Pemetrexed is infused over 10 minutes. Cisplatin is infused over one hour, with prescribed hydration before and afterwards. Blood-result review, preparation, support medicines and observation add to the visit.

STEP 3

### Access

Treatment uses suitable intravenous (IV) access. No routine take-home infusion pump is used.

STEP 4

### Home support

Take the prescribed folic acid, vitamin B12 and rash-prevention steroid plan; continue folic acid for 21 days after the final pemetrexed dose. Cisplatin needs hydration, electrolyte checks and monitoring of kidneys, hearing and nerves.

STEP 5

### Surgical review

Keep the oncology and surgical teams informed about new symptoms and other appointments.

Sources: NCCP 00849 v2 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.