BASELINE → TREATMENT → FOLLOW-UP

Tests before, during and after treatment

National Cancer Control Programme (NCCP) specifies regular blood/urine/thyroid tests but no universal frequency for each. Ask your team to write the dates. Scans and tumour markers are individualised.

Check	Before starting	First month	Later cycles	A hold or procedure	After treatment
Blood pressure	Control existing hypertension	Weekly blood-pressure checks	Monthly and as clinically indicated	Review control before restarting	Follow persistent hypertension as agreed
Blood count, kidneys, liver and thyroid	Baseline tests	Regular tests on the team plan	Continue regular tests	Repeat abnormal tests as directed	Follow unresolved effects
Urine protein	Baseline urine dipstick	Regular urine checks	Extra measurement if protein is detected	Review kidney/urine findings before restarting	Individual follow-up if abnormal
international normalised ratio (blood-clotting test) (INR) if taking warfarin	Review anticoagulation	Weekly INR until a stable dose	Then before each cycle	Additional checks when medicines or health change	Anticoagulation-team plan
Scans and clinical review	Confirm previous treatments and goals	Review symptoms, skin, nutrition and adherence	Disease assessment as directed by consultant	Team authorises any restart	Agree subsequent care and contacts

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00890 v2a official fruquintinib regimen

Tests in your plan

Carcinoembryonic antigen (CEA) — colorectal plans

If part of your plan

WHEN

For a colorectal cancer plan, if your team uses CEA; your team sets the interval.

A marker trend interpreted alongside symptoms, examination and scans.

Record the baseline, latest result and next date. A marker change alone does not establish progression.

THE WHOLE TREATMENT JOURNEY

Fruquintinib (Fruzaqla) monotherapy: treatment and checks

Metastatic colorectal cancer after standard therapies and trifluridine/tipiracil or regorafenib

Take the prescribed fruquintinib dose once daily on Days 1–21, then take no fruquintinib on Days 22–28. Take it at about the same time each day, with or without food. Repeat only when your team authorises the next cycle.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First month	Later cycles	A hold or procedure	After treatment
Blood pressure NCCP 00890	Control existing hypertension	Weekly blood-pressure checks	Monthly and as clinically indicated	Review control before restarting	Follow persistent hypertension as agreed
Blood count, kidneys, liver and thyroid NCCP 00890	Baseline tests	Regular tests on the team plan	Continue regular tests	Repeat abnormal tests as directed	Follow unresolved effects

Care / check	Before starting	First month	Later cycles	A hold or procedure	After treatment
Urine protein NCCP 00890	Baseline urine dipstick	Regular urine checks	Extra measurement if protein is detected	Review kidney/urine findings before restarting	Individual follow-up if abnormal
INR if taking warfarin NCCP 00890	Review anticoagulation	Weekly INR until a stable dose	Then before each cycle	Additional checks when medicines or health change	Anticoagulation-team plan
Scans and clinical review NCCP 00890	Confirm previous treatments and goals	Review symptoms, skin, nutrition and adherence	Disease assessment as directed by consultant	Team authorises any restart	Agree subsequent care and contacts

What each check is for

1

21 capsule days, then seven rest days

28-day cycle

Fruquintinib is a targeted capsule treatment to help control bowel cancer that has spread after previous treatments. The team checks which treatments you have received, whether they remain suitable, and your priorities.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00890

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-05 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.35