Adjuvant ribociclib + endocrine therapy

The usual adjuvant starting dose is ribociclib 400 mg once daily on Days 1–21, followed by Days 22–28 off, for up to three years. The aromatase inhibitor continues every day on its own plan; ovarian suppression is added when indicated.

BASELINE → TREATMENT → FOLLOW-UP

Treatment and monitoring over time

Columns show sequence and general intervals, not equal periods or booked appointments. The team confirms who arranges each test, who reviews results and who authorises treatment or restart.

Check	Before first dose	Cycles 1–2	Cycles 3–6	Later treatment	Completion / change
Ribociclib and endocrine partner	Confirm adjuvant 400 mg / aromatase inhibitor (AI) plan	Days 1–21 on; Days 22–28 off	Repeat if cleared; endocrine partner continues	Up to three years	Review endocrine continuation separately
Blood counts / renal / liver	Baseline profiles	Every two weeks in both cycles	Before each cycle	As clinically indicated; more often if abnormal	Follow abnormalities before restart or after stopping
electrocardiogram (ECG) / heart rhythm	Baseline ECG; check QTcF eligibility	Repeat around cycle 1 Day 14	Further ECG when clinically indicated	Repeat for symptoms, QT change or interacting medicines	No automatic restart after a QT-related hold
Electrolytes	Potassium, calcium, magnesium and phosphorus	Before each cycle; correct abnormalities	Before each cycle through cycle 6	As clinically indicated	Review unresolved abnormalities
Bone / endocrine health	Assess AI/ovarian function suppression (OFS) bone and menopausal risks	Support symptoms and adherence	AI/OFS and any injections follow their own plan	Repeat bone assessment according to risk	Longer endocrine and bone plan continues
Early-breast follow-up	Confirm stage and prior local treatment	Assess new symptoms promptly	Follow early-breast surveillance plan	Mammography/follow-up as agreed; no automatic metastatic scan schedule	Continue surveillance

THE WHOLE TREATMENT JOURNEY

Adjuvant ribociclib + endocrine therapy: treatment and checks

Adjuvant · HR-positive / HER2-negative early breast cancer

The usual adjuvant starting dose is ribociclib 400 mg once daily on Days 1–21, followed by Days 22–28 off, for up to three years. The aromatase inhibitor continues every day on its own plan; ovarian suppression is added when indicated.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before first dose	Cycles 1–2	Cycles 3–6	Later treatment	Completion / change
Ribociclib and endocrine partner NCCP 00892 · SMPC	Confirm adjuvant 400 mg / AI plan	Days 1–21 on; Days 22–28 off	Repeat if cleared; endocrine partner continues	Up to three years	Review endocrine continuation separately
Blood counts / renal / liver NCCP 00892 · SMPC	Baseline profiles	Every two weeks in both cycles	Before each cycle	As clinically indicated; more often if abnormal	Follow abnormalities before restart or after stopping
ECG / heart rhythm NCCP 00892 · SMPC	Baseline ECG; check QTcF eligibility	Repeat around cycle 1 Day 14	Further ECG when clinically indicated	Repeat for symptoms, QT change or interacting medicines	No automatic restart after a QT-related hold
Electrolytes NCCP 00892 · SMPC	Potassium, calcium, magnesium and phosphorus	Before each cycle; correct abnormalities	Before each cycle through cycle 6	As clinically indicated	Review unresolved abnormalities
Bone / endocrine health NCCP 00892 · SMPC	Assess AI/OFS bone and menopausal risks	Support symptoms and adherence	AI/OFS and any injections follow their own plan	Repeat bone assessment according to risk	Longer endocrine and bone plan continues
Early-breast follow-up NCCP 00892 · SMPC	Confirm stage and prior local treatment	Assess new symptoms promptly	Follow early-breast surveillance plan	Mammography/follow-up as agreed; no automatic metastatic scan schedule	Continue surveillance

What each check is for

1 Before the first dose

Before first dose

Sources: NCCP 00892 · SMPC

2 Early toxicity checks

Cycles 1–2

Sources: NCCP 00892 · SMPC

3 Benefit, tolerability and adherence

Later treatment

Sources: NCCP 00892 · SMPC

4 Completion or treatment change

Completion / change

Sources: NCCP 00892

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

SACTCheck · Developed by Dr Paul O'Brien · v0.84.21 · Sources checked 2026-10-25

Source reconciled; independent clinical, pharmacy and patient/PPI review pending