Serplulimab, carboplatin and etoposide

Induction lasts four planned 21-day cycles unless treatment needs to stop earlier. On Day 1 you receive serplulimab, carboplatin and etoposide by drip. With the standard intravenous (IV) schedule, etoposide is repeated on Days 2 and 3. After induction, carboplatin and etoposide stop. Serplulimab alone can continue every three weeks while it is helping and its risks remain acceptable. There is no fixed universal number of maintenance cycles in this NCCP regimen. Your team may instead prescribe etoposide by drip on Day 1 and capsules twice daily on Days 2 and 3. This is a separate alternative, not extra treatment. Oral and IV doses are not interchangeable milligram for milligram. Confirm which plan you have; take the capsules on an empty stomach as prescribed.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hormones / nervous system

Serplulimab

Severe weakness, headache, dizziness, thirst, confusion or seizures need prompt or emergency assessment.

3 Heart

Serplulimab

Chest pain, racing heart or collapse need urgent assessment.

5 Kidneys


Carboplatin / serplulimab

Much less urine, swelling or persistent vomiting need advice; kidney function affects doses.

7 Skin / nerves

Serplulimab / chemotherapy

Blisters, peeling, mouth or eye sores, new weakness or numbness need advice.



2 Lungs

Serplulimab

New cough or breathlessness: contact the team; severe breathing difficulty needs emergency help.

4 Liver

Serplulimab / etoposide

Yellow skin or eyes, dark urine or new severe nausea need advice; tests can detect changes before symptoms.

6 Bowel / pancreas / mouth

Serplulimab / chemotherapy

New diarrhoea, severe abdominal pain, blood in stool or inability to drink need prompt assessment.

8 Blood / infection

Carboplatin / etoposide; rarely immune effects

Fever, shivering or sudden illness: call oncology immediately. Report unusual bleeding or bruising.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Serplulimab

- Immune inflammation of lungs, bowel, liver, kidneys, skin and hormone glands
- Immune effects can also involve the heart, pancreas, nervous system or blood cells
- Severe infusion reactions
- Some hormone effects need lasting replacement treatment; immune effects can begin after the final dose

Carboplatin

- Low blood counts, infection or bleeding
- Nausea and vomiting
- Allergic reactions
- Kidney, hearing and nerve effects

Etoposide

- Low blood counts, infection and bleeding
- Hair loss and mouth or bowel symptoms
- Infusion reactions and low blood pressure
- A delayed risk of treatment-related leukaemia

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Induction: plan each treatment day

Day 1 carboplatin has a stronger anti-sickness plan. Days 2–3 etoposide also need planned cover, with review for two days after the final chemotherapy dose. Ask for one reconciled schedule; do not add another steroid or anti-sickness course yourself.

Maintenance: a different prescription

Serplulimab alone usually does not need the routine chemotherapy anti-sickness course. Continue the alert card and immune symptom checks; new nausea or vomiting may need investigation, not just extra anti-sickness tablets.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00908/

Sources and review status are available online. Follow your own prescription and hospital instructions.