INFORMATION TO KEEP WITH YOU

# My complete treatment handbook

## Tucatinib + capecitabine + trastuzumab

Understand your treatment. Know what to expect. Keep your plan to hand.

**Displayed calendar**
**21 days**
See the active treatment phase

**Treatment stages**
**1**
Explained in the treatment reference

**Regimen source**
**NCCP 00936**
Version 1

Tucatinib twice daily every day; capecitabine twice daily on Days 1–14 then off on Days 15–21; trastuzumab every 21 days by intravenous (IV) infusion or a locally approved under-the-skin injection. Continue while beneficial and tolerated, with clinician-approved changes.

## Treatment stages

01 **Three-medicine treatment**
Repeated 21-day cycles

Your team confirms the active stage and dates. The full treatment guide explains each step.

## Inside your handbook


| | | PAGE |
|---|---|---|
| 01 | Understanding your treatment | 2 |
| 02 | Your schedule, tests and appointments | 5 |
| 03 | Symptoms and supportive care | 12 |
| 04 | Your record and contact card | 16 |
| 05 | Sources and further information | 19 |


**My name** _____

**Hospital / oncology team** _____

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

SACTCHECK · CHAPTER 01

# Understanding your treatment

Tucatinib + capecitabine + trastuzumab · National Cancer Control Programme (NCCP) 00936

## When to get help

**Emergency: call 112 / 999**

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

**Contact your oncology service immediately**

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Report new breathlessness, jaundice, marked bleeding, severe diarrhoea, inability to drink, chest symptoms or a severe rash promptly.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.


## Your medicines and their different roles

The three medicines work together to control previously treated HER2-positive advanced breast cancer. Treatment is reviewed regularly for benefit, symptoms and side effects.

HER2 TYROSINE KINASE INHIBITOR

### Tucatinib

Tukysa · oral targeted treatment



**Blocks HER2 signals inside cells.**

Tucatinib reduces signalling from the internal part of HER2. It works alongside trastuzumab and capecitabine, which act in different ways. Its tablet schedule continues through the capecitabine break.

Reach inside the cell → Block HER2 signalling → Reduce growth signals


**What to watch for**

Report diarrhoea early, especially with poor intake. Liver tests and a full medicines review are important.

ORAL CHEMOTHERAPY

## Capecitabine

Xeloda



### Becomes fluorouracil in the body

The body converts capecitabine into fluorouracil, which disrupts deoxyribonucleic acid (the cell's genetic material) (DNA) and RNA processes needed for cell growth. Tablets are active chemotherapy; taking them at home still requires a clear monitoring and interruption plan.

Swallowed tablet → Converted to fluorouracil → Cell growth disrupted

### What to watch for

Stop tablets and contact the team for significant diarrhoea, painful mouth ulcers, hand-foot soreness affecting activities, fever or dehydration. Chest pain needs urgent assessment. Do not restart until instructed.

HER2-TARGETED ANTIBODY

**Trastuzumab**



### Targets HER2 growth signals

Trastuzumab binds to the HER2 protein on cells and helps reduce HER2-driven growth and recruit immune activity. HER2 testing identifies the cancers for which this approach is used.

HER2 on cell surface → Antibody binds → Growth signalling reduced

### What to watch for

Attend planned echoes. Report breathlessness, ankle swelling, rapid weight gain or infusion symptoms.

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

## Your treatment aim

### Indication

Advanced / metastatic · previously treated HER2-positive breast cancer

### Where this fits

Tucatinib twice daily every day; capecitabine twice daily on Days 1–14 then off on Days 15–21; trastuzumab every 21 days by the agreed intravenous (IV) or locally approved subcutaneous (SC) route. Continue while beneficial and tolerated, with clinician-approved changes.

## HER2CLIMB · longer follow-up

This explains the study setting and its meaning. Ask the team how the evidence applies to you.

Adding tucatinib helped delay cancer growth and improved survival compared with the other study group. The study does not predict an individual outcome or compare this combination with every newer treatment.

SACTCHECK · CHAPTER 02


# Your schedule, tests and appointments

Tucatinib + capecitabine + trastuzumab · NCCP 00936

## Each phase in your treatment plan

**Tucatinib + capecitabine + trastuzumab**

21-day view




This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

REPEATED 21-DAY CYCLES

### Three-medicine treatment

**Repeated 21-day cycles**

The capecitabine break does not mean a tucatinib break. Your team may hold or adjust one medicine without applying the same change to all three. Trastuzumab route is agreed separately; the fixed subcutaneous (SC) dose must not be treated as an intravenous (IV) loading or maintenance dose.

The order depends on the indication and your recovery. The whole-treatment map shows how local treatment and ongoing medicines relate to each other.

BASELINE → TREATMENT → FOLLOW-UP

## When checks happen

Elapsed-time tests remain due if a cycle is delayed. The treatment team books, reviews and acts on results. Calendar timing alone does not authorise a dose.

| Check | Before starting | Every 3-week cycle | Around week 12 | Ongoing treatment | Treatment held / changed |
|---|---|---|---|---|---|
| **Blood count / kidney function** | full blood count (FBC), renal profile and creatinine clearance | FBC and renal profile before each cycle; sooner for diarrhoea, dehydration or infection | Continue each-cycle tests | Interpret creatinine changes with hydration and true renal function; tucatinib can affect creatinine transport | Reassess renal function and capecitabine safety before restarting |

| Check | Before starting | Every 3-week cycle | Around week 12 | Ongoing treatment | Treatment held / changed |
|---|---|---|---|---|---|
| **Liver tests** | alanine aminotransferase (ALT), aspartate aminotransferase (AST) and bilirubin | Every 3 weeks, or sooner if symptoms / abnormalities | Continue every 3 weeks | Report jaundice or dark urine promptly | Repeat abnormal tests and review tucatinib/ capecitabine separately |
| **dihydropyrimidine dehydrogenase (a medicine-processing enzyme) (DPD) / capecitabine safety** | Confirm pretreatment DPD result before first capecitabine; renal impairment can affect uracil phenotyping | Review early severe diarrhoea, mouth ulcers, infection or neurological symptoms immediately | No routine repeat DPD test simply because a cycle passes | Complete deficiency excludes capecitabine; partial deficiency requires an individual plan | Hold affected tablets and seek urgent advice; do not restart independently |
| **Heart function** | Cardiac history, electrocardiogram (ECG) and baseline echo/ multigated acquisition (heart pumping) scan (MUGA) under trastuzumab product information | Review breathlessness, swelling and chest pain at each visit | Repeat left ventricular ejection fraction (heart pumping measurement) (LVEF) approximately 12-weekly under the NCCP/ individual cardiac plan | 4–8-weekly if impairment; keep elapsed-time due dates when route changes | A qualifying LVEF decline requires hold and repeat within 3 weeks; post-treatment follow-up is planned separately |
| **Diarrhoea / skin / mouth** | Provide separate tablet-hold instructions and hydration plan | Review bowels, intake, hands/ feet and mouth before continuation | Symptoms can occur between visits and during capecitabine breaks | Tucatinib continues on the planned Days 15–21 capecitabine break unless specifically held | The medicines have different interruption and restart rules |

| Check | Before starting | Every 3-week cycle | Around week 12 | Ongoing treatment | Treatment held / changed |
|---|---|---|---|---|---|
| **Disease / central nervous system (CNS) review** | Baseline disease assessment and CNS plan when indicated | Clinical review; imaging not tied to tablet-break days | Response assessment commonly every 2–4 months, individualised | Earlier assessment for new symptoms; CNS schedule decided separately | Markers support clinical/ imaging assessment, not decisions alone |

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00936 v1 · official regimen · ESC 2022 cardio-oncology guideline · cardiac surveillance and risk · ASCO · biomarkers in metastatic breast cancer · ESMO metastatic breast cancer guideline · response assessment · EMA Tukysa product information · retrieved 25 October 2026 · EMA Xeloda product information · retrieved 25 October 2026 · EMA Herceptin product information · retrieved 25 October 2026

THE WHOLE TREATMENT JOURNEY

## Tucatinib + capecitabine + trastuzumab: treatment and checks

Tucatinib twice daily every day; capecitabine twice daily on Days 1–14 then off on Days 15–21; trastuzumab every 21 days by IV infusion or a locally approved under-the-skin injection. Continue while beneficial and tolerated, with clinician-approved changes.

**Read from left to right.** Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

**Treatment, tests and reviews across the course**

| Care / check | Before starting | Days 1–14 | Days 15–21 | Next cycle | Disease review |
|---|---|---|---|---|---|
| **Cancer medicines** NCCP 00936 | Confirm prior HER2 therapies and DPD result | Tucatinib + capecitabine; trastuzumab Day 1 by agreed IV or SC formulation | Tucatinib continues; capecitabine break | Review before any restart / next infusion | Continue, modify or change treatment according to benefit / toxicity |
| **Surgery** METASTATIC | No routine curative surgery sequence from this regimen | Local procedures only if indicated | Coordinate interruptions with multidisciplinary team (MDT) | Review symptoms and functional needs | Site-specific / CNS MDT when needed |
| **Radiotherapy** METASTATIC | Assess need for symptom or brain-directed treatment | Any concurrent radiotherapy (RT) needs an explicit plan | Do not assume capecitabine can continue with RT | Coordinate holds / restart with radiation team | Imaging and local treatment depend on disease site |

| Care / check | Before starting | Days 1–14 | Days 15–21 | Next cycle | Disease review |
|---|---|---|---|---|---|
| **Blood tests**<br>NCCP 00936 | FBC, renal, liver; DPD testing before capecitabine | Extra tests if symptoms / toxicity | Pre-cycle bloods and symptom review | FBC / renal per cycle; liver at least every 3 weeks | Check dehydration, hepatic toxicity and interactions |
| **Heart checks**<br>NCCP 00936 · HERCEPTIN · ESC | Baseline cardiac assessment, ECG and echo/MUGA | Report chest pain or breathlessness | Symptoms need assessment between visits | Follow the recorded LVEF plan, approximately every 12 weeks | Risk / abnormalities may shorten interval |
| **Cancer imaging**<br>METASTATIC · MARKERS | Baseline disease sites; brain assessment as indicated | No scan required simply because Day 14 is reached | Imaging schedule independent of tablet break | Usually review response about every 2–4 months; individualise | Use consistent modalities; earlier imaging if concern |

## What each check is for

1

### Blood tests and treatment review

FBC, renal, liver; DPD testing before capecitabine → Extra tests if symptoms / toxicity → Pre-cycle bloods and symptom review → FBC / renal per cycle; liver at least every 3 weeks → Check dehydration, hepatic toxicity and interactions

Results may change whether a dose is given or held.

**Ask / act:** Confirm the booking and who reviews the result before treatment.

2

### Heart tests

Baseline echo / LVEF → Report chest pain or breathlessness → Symptoms need assessment between visits → Repeat echo according to HER2 plan, commonly 12-weekly → Risk / abnormalities may shorten interval

ECG measures rhythm; echo measures pumping function. They are different checks.

**Ask / act:** Keep the planned appointments and report symptoms between them.

3

### Imaging and markers

Baseline disease sites; brain assessment as indicated → No scan required simply because Day 14 is reached → Imaging schedule independent of tablet break → Usually review response about every 2–4 months; individualise → Use consistent modalities; earlier imaging if concern

The purpose and frequency depend on early versus advanced disease.

**Ask / act:** Ask which test is next and what decision its result will inform.

4

### Who arranges and reviews each check?

At initiation and every handover

A recommended test can be missed if responsibility is unclear.

**Ask / act:** Ask the team to document the booking owner, result reviewer, due window and escalation route.

5

### Trastuzumab formulation confirmed

Before first dose and any route change

IV and SC doses and missed-dose instructions differ.

**Ask / act:** Confirm the route and visit/observation plan with the team.

### At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

## Your practical treatment plan

STEP 1

### Before dispensing

Confirm blood tests, interactions, the exact tablet strengths and the plan for missed doses.

STEP 2

### Taking tablets

STEP 3

### Review before restarting

The team confirms blood results, side effects and the next prescription. Never double a missed dose or restart held treatment without advice.

STEP 4

### Trastuzumab visit

Trastuzumab every three weeks with staff observation. IV treatment may use a cannula, peripherally inserted central catheter (PICC) or port; the locally approved SC alternative uses an injection under the skin; no routine take-home infusion pump.

STEP 5

### Confirm the trastuzumab route

Your team confirms IV infusion or the locally approved under-the-skin injection. The visit length, observation and missed-dose instructions differ; ask before switching. The injection has no loading dose.

### Missed or vomited dose

Do not take an extra dose. Follow the medicine-specific instructions and ask the team if uncertain.

### Safe handling

Swallow tablets whole, keep them in their original packaging away from children, and return unused tablets to pharmacy.

### Practical arrangements

Confirm who supplies the tablets, who arranges tests, who reviews results and your urgent contact route.

### Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

### PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

## Before starting and between visits

Plan the tests, access and support before the first dose.

### Blood tests

FBC, renal, liver; DPD testing before capecitabine → Extra tests if symptoms / toxicity → Pre-cycle bloods and symptom review → FBC / renal per cycle; liver at least every 3 weeks → Check dehydration, hepatic toxicity and interactions

### Heart checks

Baseline echo / LVEF → Report chest pain or breathlessness → Symptoms need assessment between visits → Repeat echo according to HER2 plan, commonly 12-weekly → Risk / abnormalities may shorten interval

## Imaging, local treatment and follow-up

The whole pathway has several connected parts.

### Scans and imaging

Baseline disease sites; brain assessment as indicated → No scan required simply because Day 14 is reached → Imaging schedule independent of tablet break → Usually review response about every 2–4 months; individualise → Use consistent modalities; earlier imaging if concern

### Radiotherapy relationship

Assess need for symptom or brain-directed treatment → Any concurrent RT needs an explicit plan → Do not assume capecitabine can continue with RT → Coordinate holds / restart with radiation team → Imaging and local treatment depend on disease site

### Bone health

If an aromatase inhibitor or ovarian suppression is prescribed, assess fracture risk and baseline dual-energy X-ray absorptiometry (bone density) (DEXA) as indicated. Repeat timing depends on the result, risk and bone treatment; it is not an automatic scan for everyone.

# Symptoms and supportive care

Tucatinib + capecitabine + trastuzumab · NCCP 00936

## Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

| Treatment or check | Timing in this view |
|---|---|
| **Tucatinib** |  **Days 1–21** Twice daily, continuous |
| **Capecitabine ON** |  **Days 1–14** Twice daily |
| **Capecitabine OFF** |  **Days 15–21** 7-day break |
| **Trastuzumab** |  **Day 1** IV or agreed SC formulation |
| **Diarrhoea / hand-foot** |  **Days 1–21** Report early, including during break |
| **Pre-cycle review** |  **Days 19–21** Timing arranged by unit |

**Treatment bands:** Show the planned medicine schedule, subject to review and dose changes.

**Checks and restart:** Keep the arranged blood / heart appointments. A calendar day alone does not authorise the next dose.

**Symptoms:** Can occur at any time; broad bands are not measured risk curves.

> **Why these bands stay broad:** Planned treatment days are distinct from broad possible symptom patterns. Symptoms can occur outside these bands. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00936 v1 · official regimen ↗

**There are no guaranteed "safe days".** Fever, shivers, sudden illness, new breathing problems or severe symptoms need prompt advice on any day, including a treatment break.

### Between-treatment notes

### Day 1–21: Your reference cycle

Tucatinib twice daily every day; capecitabine twice daily on Days 1–14 then off on Days 15–21; trastuzumab every 21 days by IV infusion or a locally approved under-the-skin injection. Continue while beneficial and tolerated, with clinician-approved changes. Check the agreed prescription and contact the team for new symptoms. Restart only after the team confirms the plan.

## Support around the medicines

**1** **Support for this phase**

Obtain a clear diarrhoea action plan and separate instructions for each tablet. Anti-sickness medicines and interactions require individual review.

**2** **Bloods and cardiac checks**

FBC, renal, liver; DPD testing before capecitabine → Extra tests if symptoms / toxicity → Pre-cycle bloods and symptom review. Symptoms need assessment between visits.

**3** **Visits and practical support**

Ask about travel, work, fertility, menopause, exercise and help at home. The team arranges and reviews clinical checks; the patient should not have to calculate the monitoring timetable.

## Questions worth asking

### When are my next tests?

Your team should confirm the next test, who books it, and who reviews the result before treatment. The map shows expected intervals; it does not track completed tests.

### Does every scan follow a fixed schedule?

Use consistent modalities; earlier imaging if concern. New symptoms can require earlier assessment.

**Try explaining the plan back:** Which medicines are you taking now? What changes in the next phase? Which checks are due and who reviews them before treatment?

**Tucatinib**

- Diarrhoea and dehydration
- Liver test changes or liver injury
- Nausea and fatigue
- Clinically important medicine interactions

### Capecitabine

- Hand-foot syndrome
- Mouth ulcers / reduced intake
- Low blood counts and infection
- Cardiotoxicity including chest pain / ischaemia
- Severe early toxicity including dihydropyrimidine dehydrogenase (a medicine-processing enzyme) (DPD)-related toxicity
- Renal impairment and medicine interactions

### Trastuzumab

- Heart pumping-function reduction / heart failure
- Infusion or allergic reaction
- Diarrhoea and fatigue
- Rare serious lung effects

---

**Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Tucatinib + capecitabine + trastuzumab

**Active medicines:** Tucatinib + Capecitabine + Trastuzumab

**Emetogenic risk:** Tucatinib / capecitabine oral minimal to low; trastuzumab minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

### Additional phase-specific support

- Agree a personalised antiemetic and diarrhoea prescription; review CYP and other interactions. No repeated intravenous (IV) chemotherapy premedication is assumed for these tablets.
- Tucatinib continues every day; capecitabine is Days 1–14 then off Days 15–21. Confirm separate hold / restart instructions.
- Premedicines are not usually required. If prior hypersensitivity, consider paracetamol / antihistamine under the clinician plan.
- Explain when to hold capecitabine and call for diarrhoea, mouth soreness or painful hand-foot symptoms. Assess dehydration promptly.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00936 · VERSION 1

# Tucatinib + capecitabine + trastuzumab

Tucatinib twice daily every day; capecitabine twice daily on Days 1–14 then off on Days 15–21; trastuzumab every 21 days by IV infusion or a locally approved under-the-skin injection. Continue while beneficial and tolerated, with clinician-approved changes.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

## A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

### 1 Diarrhoea / dehydration

**Tucatinib + capecitabine**

Report diarrhoea early. Follow the separate tablet-hold instructions; reduced urine or dizziness needs prompt advice.

### 3 Liver injury


**Tucatinib**

Keep three-weekly liver blood tests. Report yellow skin / eyes or dark urine.

### 5 Blood counts / infection

**Capecitabine**

Fever, shivers or sudden illness: call promptly. Blood tests guide the next cycle.




### 2 Hands / feet

**Capecitabine**

Painful redness or peeling affecting activity: stop capecitabine and contact oncology.

### 4 Heart / chest symptoms

**Trastuzumab / capecitabine**

Keep echoes. New chest pain: stop capecitabine and seek urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

SACTCHECK · CHAPTER 04

# Your record and contact card

Tucatinib + capecitabine + trastuzumab · National Cancer Control Programme (NCCP) 00936

Record each cycle separately. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Use a new record sheet for each cycle or review. Copy or print another sheet when needed; the number of sheets does not prescribe the length of treatment.

Cycle number: _____ Start date: _____ Next review: _____

**Three-medicine treatment · reusable record** · copy this sheet for another cycle of the same phase.

## Tucatinib · HER2 tyrosine kinase inhibitor

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/ day or at night: call | □ 7+ extra stools/ day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Nausea** | □ None | □ Queasy but eating and drinking | □ Intake reduced: contact the team | □ Unable to maintain intake: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

## Capecitabine · Oral chemotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/ day or at night: call | □ 7+ extra stools/ day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Hand-foot soreness** | □ None | □ Redness or tingling without pain | □ Pain limits usual tasks: call | □ Pain limits washing / dressing: urgent review | □ Acute severe illness: emergency help |
| **Mouth soreness / ulcers** | □ None | □ Mild soreness; eating normally | □ Painful; soft food needed: call | □ Cannot eat / drink enough: urgent review | □ Major breathing difficulty: 112/999 |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

## Trastuzumab · HER2-targeted antibody

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/ day or at night: call | □ 7+ extra stools/ day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Carry your treatment details

Print, cut out and fold. Follow the contact plan from your own hospital.

SACTCHECK · SACT TREATMENT

## Tucatinib + capecitabine + trastuzumab

NCCP 00936 v1 · 21-day cycle

Tucatinib twice daily every day; capecitabine twice daily on Days 1–14 then off on Days 15–21; trastuzumab every 21 days by IV infusion or a locally approved under-the-skin injection. Continue while beneficial and tolerated, with clinician-approved changes.

| 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|

**Your contact plan**

Oncology team: Insert numbers here
Out of hours: Insert numbers here.

Evenings / overnight: instructed out-of-hours pathway or nearest ED.
**Emergency: 112 / 999.**

Other hospital / own contact: _____

QR: https://sactcheck.com/patient/00936/ · no patient information.

Draft for review · SACTCheck v0.84.21

SACTCHECK · CHAPTER 05

# Sources and further information

Tucatinib + capecitabine + trastuzumab · National Cancer Control Programme (NCCP) 00936

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.


1. **NCCP 00936 v1 · official regimen ↗**
   https://healthservice.hse.ie/documents/8391/936_Tucatinib_Regimen.pdf
2. **ESC 2022 cardio-oncology guideline · cardiac surveillance and risk**
   https://academic.oup.com/eurheartj/article/43/41/4229/6673995
3. **ASCO · biomarkers in metastatic breast cancer**
   https://ascopubs.org/doi/10.1200/JCO.2015.61.1459
4. **ESMO metastatic breast cancer guideline · response assessment**
   https://www.annalsofoncology.org/article/S0923-7534(26)00902-6/fulltext
5. **EMA Tukysa product information · retrieved 25 October 2026**
   https://www.ema.europa.eu/en/documents/product-information/tukysa-epar-product-information_en.pdf
6. **EMA Xeloda product information · retrieved 25 October 2026**
   https://www.ema.europa.eu/en/documents/product-information/xeloda-epar-product-information_en.pdf
7. **EMA Herceptin product information · retrieved 25 October 2026**
   https://www.ema.europa.eu/en/documents/product-information/herceptin-epar-product-information_en.pdf
8. **eviQ · implanted port patient information**
   https://www.eviq.org.au/p/507
9. **ESMO 2024 early breast cancer guideline · treatment and follow-up**
   https://www.annalsofoncology.org/article/S0923-7534(23)05104-9/fulltext
10. **HSE · chemotherapy side effects and urgent advice**
    https://www2.hse.ie/conditions/chemotherapy/side-effects/
11. **NICE NG101 · bone health and follow-up**
    https://www.nice.org.uk/guidance/ng101/chapter/Recommendations
12. **NOGG · section 7, aromatase inhibitors and fracture risk**
    https://www.nogg.org.uk/full-guideline/section-7-strategies-management-osteoporosis-and-fracture-risk
13. **eviQ · peripherally inserted central catheter (PICC) patient information**
    https://www.eviq.org.au/p/543
14. **NCCP NCIS antiemetics v8 (2025), Table 1**
    https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf
15. **HER2CLIMB · original analysis (2020)**
    https://pubmed.ncbi.nlm.nih.gov/31825569/
16. **HER2CLIMB final overall survival analysis (2022)**
    https://pubmed.ncbi.nlm.nih.gov/34954044/