Nivolumab · every 2 weeks · previously treated non-small-cell lung cancer (NSCLC)

Aim to control advanced lung cancer, relieve symptoms and prolong life where possible.

Previously treated locally advanced or metastatic non-small-cell lung cancer (NSCLC). · NCCP 00483 v13a · Document revised 9 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

One treatment visit every 2 weeks; your team confirms duration.

14 day view

Treatment timing in this window

Treatment — Day 1

Nivolumab monotherapy

Nivolumab 240 mg intravenous (IV) over 30 minutes OR 600 mg as a subcutaneous injection over 3–5 minutes on Day 1 every 2 weeks. Your team confirms one formulation; the whole visit includes checks and preparation. Continue until progression or unacceptable toxicity, with individual benefit and safety review.

A typical 14-day cycle

Day 1	Day 2–14
Your treatment visit	Between visits

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** NCCP	Blood count, kidney/liver, glucose and thyroid tests; clinical and medicine review.
During	**Bloods and symptoms** NCCP	Blood count, kidney/liver and glucose before each cycle. Thyroid once a month. Cortisol if indicated.
Response / after	**Individual plan** NCCP / individual plan	Review progression before cycle 8. Monitor immune effects for at least 5 months after the last dose.

Between treatments

- Report new symptoms promptly between visits and after treatment stops.
- Follow the written steroid or hormone-replacement plan. Do not restart a held medicine yourself.
- If your team has permanently stopped nivolumab because of a serious immune reaction, improvement does not automatically make restarting safe. Keep that decision in your treatment record; only your specialist team can review further treatment.

Bring your medicine list, alert card, test dates and actual urgent contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hormones / nervous system

Nivolumab

Severe headache, confusion, weakness, dizziness or marked thirst need prompt advice.

3 Heart

Nivolumab

Chest pain, racing heartbeat or fainting needs urgent assessment.

5 Kidneys


Nivolumab

Report much less urine, new swelling or a change in kidney tests.

7 Skin / muscles

Nivolumab

Spreading rash, blisters, severe weakness or muscle pain needs prompt review.



2 Lungs

Nivolumab

New cough or breathlessness can signal inflammation or another urgent cause.

4 Liver

Nivolumab

Yellow eyes/skin or dark urine may signal inflammation; tests also check the liver.

6 Bowel

Nivolumab

New or persistent diarrhoea, blood in stools or severe tummy pain needs urgent advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact the oncology team immediately for new or worsening breathlessness, chest pain, persistent diarrhoea, severe tummy pain, yellow skin/eyes, much less urine, severe headache, marked weakness, confusion, unusual thirst or passing much more urine, or a spreading/blistering rash. For severe breathing difficulty, collapse, seizure or other immediately dangerous symptoms, seek emergency care. Do not wait for the next infusion.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Previously treated locally advanced or metastatic non-small-cell lung cancer (NSCLC). · NCCP 00483 v13a · Document revised 9 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Nivolumab · lung

Every 2 weeks; confirm duration and current phase.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.