Pembrolizumab 400 mg · every 6 weeks · lung cancer

For metastatic lung cancer, the aim is to control cancer and help people live longer. After surgery and platinum chemotherapy, the adjuvant aim is to reduce the chance of cancer returning. Ask which aim applies to you.

Lung cancer only: first-line metastatic non-small-cell lung cancer (NSCLC) or adjuvant NSCLC after surgery and platinum chemotherapy · NCCP 00558 v13 · Document revised 9 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

One treatment visit every 6 weeks; your team confirms duration.

42 day view

Treatment timing in this window

Infusion — Day 1

Pembrolizumab every 6 weeks

After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

A typical 42-day cycle

Day 1	Day 2–42
Your infusion visit	Between visits

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** NCCP	Blood count, kidney/liver, glucose and thyroid tests; clinical and medicine review.
During	**Bloods and symptoms** NCCP	Blood count, kidney/liver and glucose before each cycle. Thyroid function tests every 3–6 weeks. Cortisol if indicated.
Response / after	**Individual plan** NCCP / individual plan	Scans on your individual plan; report delayed immune symptoms.

Between treatments

- Report new symptoms promptly between visits and after treatment stops.
- Follow the written steroid or hormone-replacement plan. Do not restart a held medicine yourself.
- A permanent-stop decision still applies after symptoms improve; your team decides whether restarting is safe.

Bring your medicine list, alert card, test dates and actual urgent contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hormones / nervous system

Pembrolizumab

Severe headache, confusion, weakness, dizziness or marked thirst need prompt advice.

3 Heart

Pembrolizumab

Chest pain, racing heartbeat or fainting needs urgent assessment.

5 Kidneys


Pembrolizumab

Report much less urine, new swelling or a change in kidney tests.

7 Skin / muscles

Pembrolizumab

Spreading rash, blisters, severe weakness or muscle pain needs prompt review.



2 Lungs

Pembrolizumab

New cough or breathlessness can signal inflammation or another urgent cause.

4 Liver

Pembrolizumab

Yellow eyes/skin or dark urine may signal inflammation; tests also check the liver.

6 Bowel

Pembrolizumab

New or persistent diarrhoea, blood in stools or severe tummy pain needs urgent advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact the oncology team immediately for new or worsening breathlessness, chest pain, persistent diarrhoea, severe tummy pain, yellow skin/eyes, much less urine, severe headache, marked weakness, confusion, unusual thirst or passing much more urine, or a spreading/blistering rash. For severe breathing difficulty, collapse, seizure or other immediately dangerous symptoms, seek emergency care. Do not wait for the next infusion.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Lung cancer only: first-line metastatic NSCLC or adjuvant NSCLC after surgery and platinum chemotherapy · NCCP 00558 v13 · Document revised 9 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Pembrolizumab · lung

Every 6 weeks; confirm duration and current phase.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.