SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Nab-paclitaxel · every 3 weeks

Metastatic breast cancer after first-line treatment · NCCP 00230 · version 7 · 21-day cycle

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Patient name	DOB	Hospital no.	Date

Treatment context
The aim is to control the cancer and its symptoms. The team reviews benefit, side effects and your priorities before continuing.

Schedule
Nab-paclitaxel on Day 1 every 21 days. Review response after two cycles; the NCCP protocol says to discontinue if there is no response. Further treatment depends on benefit and tolerance.

1. GENERIC SACT RISKS / DISCUSSION POINTS

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Off-label indication / schedule discussed where applicable
- ☐ Supportive medicines and urgent advice discussed
- ☐ Common, serious and lasting medicine-specific risks discussed below
- ☐ Exact medicine, formulation and schedule confirmed
- ☐ Blood tests and required heart / imaging checks explained
- ☐ Pregnancy, contraception and fertility discussed where relevant
- ☐ Patient priorities and questions recorded

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Nab-paclitaxel
CHEMOTHERAPY

- ☐ Numbness / tingling or weakness, sometimes lasting
- ☐ Low blood counts, infection and sepsis
- ☐ Hair loss, fatigue and muscle / joint aches
- ☐ Infusion reactions; uncommon serious lung inflammation

Discuss the exact indication, expected benefit, alternatives, uncertainty, serious and lasting risks, contraception and fertility, venous access, monitoring and the urgent contact plan. Nab-paclitaxel is an albumin-bound formulation. It is not interchangeable with ordinary paclitaxel or with the weekly nab-paclitaxel schedule.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00230 v7.

SACTCheck v0.84.1
Developed by Dr Paul O'Brien