Eribulin · Days 1 and 15 of 28

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Eribulin · Days 1 and 15 of 28 · each treatment day

Active medicines: Eribulin

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally once	Each treatment day only Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- No routine hypersensitivity premedication is specified. Low-risk antiemetic support is reviewed separately, including QT interactions.
- This is not the licensed Day 1 / Day 8 schedule. The reason for the alternative schedule, evidence limits and off-label consent are part of the treatment discussion.
- Review interactions, QT risk and electrolytes before selecting an antiemetic; individualise rather than automatically issuing this draft.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00743 v3 · p4 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-18 · SACTCheck v0.84.1 · Regimen-level drafts; prescriber and oncology-pharmacy verification required