TRAIN-2 with Phesgo

NCCP 00790 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

Phesgo is one combined injection of pertuzumab and trastuzumab under the skin of the thigh. It is not two separate injections and is never given into a vein.

STEP 1

Before the visit

Confirm blood results, cardiac checks, active phase, previous dual HER2 dose and the consent discussion.

STEP 2

The thigh injection

Loading/reloading 1200 mg/600 mg over 8 minutes with 30 minutes observation; maintenance 600 mg/600 mg over 5 minutes with 15 minutes observation. Complete the observation before chemotherapy. Alternate thighs; avoid sore, bruised or damaged skin.

STEP 3

Chemotherapy and access

After Phesgo observation, chemotherapy is given through a cannula, PICC or port. The team explains access insertion and care. No take-home infusion pump is routinely used.

STEP 4

Leaving the unit

Confirm supportive medicines, next visits and due tests. Injection time is shorter than the whole appointment; preparation, assessment and observation still take time.

Switching / a missed appointment

The team checks the last dual HER2 dose, including IV treatment: a gap of less than 6 weeks uses the maintenance dose; 6 weeks or longer needs reloading. A first Phesgo visit after IV treatment does not automatically require a loading dose.

Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

Sources: NCCP 00790 v3 · official regimen · eviQ · implanted port patient information · eviQ · PICC patient information · EMA Phesgo product information. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.