NCCP 00790 · clinician draft

TRAIN-2 with Phesgo

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Postoperative HER2-only phase · confirm selected regimen

Active medicines: Selected postoperative HER2 medicine; separate source regimen

Emetogenic risk: Confirm the selected postoperative regimen

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Open the separately selected postoperative regimen before issuing supportive medicines. Its emetic risk, premedication and tests apply.
- Phesgo or trastuzumab alone generally needs no routine antiemetic prophylaxis. Do not apply this statement automatically to an antibody-drug conjugate.
- Pathology, including residual invasive disease, informs the next HER2 medicine and remaining treatment course.
- Continue cardiac monitoring on the selected treatment plan; surgery does not restart the clock.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00790 v3 · p1 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-21 · SACTCheck v0.84.15 · Regimen-level drafts; prescriber and oncology-pharmacy verification required