NCCP 00796 · clinician draft

Phesgo + docetaxel

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Taxane + Phesgo phase

Active medicines: Docetaxel + Pertuzumab + Trastuzumab

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- Docetaxel: dexamethasone 8 mg orally twice daily for 3 days, starting the day before each dose; at least 3 doses before infusion. Missed doses require consultant advice; any 20 mg IV replacement is a consultant decision.
- Reconcile all steroid exposure and aprepitant interactions; the antiemetic steroid reference is not an instruction to add a duplicate course.
- Consider primary G-CSF under the source and individual risk plan.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00796 v4 · p5 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-21 · SACTCheck v0.84.15 · Regimen-level drafts; prescriber and oncology-pharmacy verification required