NCCP 00858 · clinician draft

Pembrolizumab 200 mg + weekly carboplatin AUC 1.5 / paclitaxel → AC

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

After surgery · pembrolizumab alone

Active medicines: Pembrolizumab

Emetogenic risk: Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- After surgery: pembrolizumab 200 mg every 21 days for nine doses OR 400 mg every 42 days for five doses. These are alternative schedules, not consecutive courses. Progression preventing definitive surgery or unacceptable pembrolizumab toxicity changes this plan; postoperative treatment is not automatic.
- No routine cytotoxic antiemetics, taxane premedicines or chemotherapy G-CSF are carried forward.
- Continue the immune-toxicity and required blood/hormone monitoring.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00858 v3 · p9 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-20 · SACTCheck v0.84.13 · Regimen-level drafts; prescriber and oncology-pharmacy verification required