BASELINE → TREATMENT → FOLLOW-UP

Tests, symptoms and treatment decisions

Medicine-safety checks, tumour-tissue testing and response imaging answer different questions. Product-information additions are labelled. No universal blood tumour-marker panel or scan interval is implied.

Check	Before starting	During treatment	Between visits	Next treatment / response	After a pause or stop
Bloods / medicine safety	Blood count, kidney/liver tests; hearing assessment and creatinine clearance when clinically indicated; NCCP lung testing for patients at high risk. Current lomustine SmPC additionally advises baseline lung-function assessment. Current cisplatin product information advises baseline audiometry. Confirm recovery after RT and the individual hydration/electrolyte plan.	Regular blood count, kidney/liver tests, with hearing and lung assessment when indicated. Current lomustine SmPC advises weekly blood counts for at least six weeks after a dose and continuing lung assessment. Current cisplatin product information additionally advises weekly kidney/liver tests, blood counts and electrolytes through treatment, with repeat hearing testing for symptoms. Review nerves and bowel function before vincristine.	Keep additional tests even on days without an anticancer dose.	Team reviews current and previous results, symptoms and dose history.	Agree continuing checks; lomustine effects may persist.
Neurology / medicines / function	Record usual symptoms, medicines, daily function and support needs.	Review headache, seizures, speech, strength, cognition and any steroid plan.	Report new changes early; do not wait for an MRI.	Discuss treatment tolerance, priorities and the next authorised dose.	Agree rehabilitation, symptom support and contacts.
Response / appointments	Confirm exact diagnosis and baseline imaging.	Imaging and clinical review follow the specialist plan.	Record bookings, result-review contacts and questions.	Interpret scans with symptoms, treatment effects and medicines.	Agree the next imaging and clinical review.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00806 v2 · current Irish regimen · NCI: medulloblastoma diagnosis, risk groups and treatment · patient education · HPRA: cisplatin Accord current product information · July 2026 · Lomustine medac current product information · January 2026

THE WHOLE TREATMENT JOURNEY

Cisplatin, lomustine and vincristine (CLV): treatment and checks

NCCP 00806: selected adults with high-risk medulloblastoma or another eligible primitive neuro-ectodermal tumour, including pinealoblastoma.

Start four to six weeks after craniospinal radiotherapy, once cleared by your team. In each 42-day cycle, lomustine is taken once on Day 1, cisplatin is infused on Day 1, and vincristine is infused on Days 1, 8 and 15. Eight cycles are planned, with individual safety review.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Next treatment / response	After a pause or stop
Bloods / medicine safety NCCP 00806	Blood count, kidney/liver tests; hearing assessment and creatinine clearance when clinically indicated; NCCP lung testing for patients at high risk. Current lomustine SmPC additionally advises baseline lung-function assessment. Current cisplatin product information advises baseline audiometry. Confirm recovery after RT and the individual hydration/electrolyte plan.	Regular blood count, kidney/liver tests, with hearing and lung assessment when indicated. Current lomustine SmPC advises weekly blood counts for at least six weeks after a dose and continuing lung assessment. Current cisplatin product information additionally advises weekly kidney/liver tests, blood counts and electrolytes through treatment, with repeat hearing testing for symptoms. Review nerves and bowel function before vincristine.	Keep additional tests even on days without an anticancer dose.	Team reviews current and previous results, symptoms and dose history.	Agree continuing checks; lomustine effects may persist.
Neurology / medicines / function NCCP 00806	Record usual symptoms, medicines, daily function and support needs.	Review headache, seizures, speech, strength, cognition and any steroid plan.	Report new changes early; do not wait for an MRI.	Discuss treatment tolerance, priorities and the next authorised dose.	Agree rehabilitation, symptom support and contacts.
Response / appointments NCCP 00806	Confirm exact diagnosis and baseline imaging.	Imaging and clinical review follow the specialist plan.	Record bookings, result-review contacts and questions.	Interpret scans with symptoms, treatment effects and medicines.	Agree the next imaging and clinical review.

What each check is for

1 Recovery after craniospinal radiotherapy

Before CLV

CLV combines three chemotherapy medicines after craniospinal radiotherapy for selected adults with high-risk medulloblastoma or another eligible embryonal tumour. It is part of a specialist treatment plan; it is not a glioma regimen.

Ask / act: Confirm the current phase, prescription, blood tests and support medicines.

Sources: NCCP 00806

2 Cisplatin, lomustine and vincristine (CLV)

42-day cycle

CLV combines three chemotherapy medicines after craniospinal radiotherapy for selected adults with high-risk medulloblastoma or another eligible embryonal tumour. It is part of a specialist treatment plan; it is not a glioma regimen.

Ask / act: Confirm the current phase, prescription, blood tests and support medicines.

Sources: NCCP 00806

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-05 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.39