SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Weekly paclitaxel

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Shared advanced-cancer protocol · NCCP 00226 · version 9 · 28-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
This weekly paclitaxel protocol is used in specified advanced breast, ovarian, small-cell lung and bladder-cancer settings. The aim is usually disease control and symptom relief. Evidence and alternatives differ between cancers; confirm the indication with your oncologist.

Schedule
Paclitaxel on Days 1, 8, 15 and 22 of a 28-day cycle; a Days 1, 8 and 15 schedule may be selected by the consultant. Confirm your appointment plan.

1. GENERIC SACT RISKS / DISCUSSION POINTS

- ☐ Purpose, expected benefit and treatment plan explained
- ☐ Treatment may be delayed, modified or stopped because of toxicity
- ☐ Unexpected, rare or serious adverse effects can occur
- ☐ Supportive medicines and urgent contact advice discussed
- ☐ Pregnancy avoidance / contraception / fertility discussed where relevant
- ☐ Bone-marrow suppression / infection risk
- ☐ Anaemia / fatigue and low platelets / bleeding
- ☐ Nausea / vomiting and bowel disturbance
- ☐ Mouth soreness / mucositis and skin changes
- ☐ Allergic / infusion reaction and other organ-specific toxicity

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Paclitaxel

TAXANE CHEMOTHERAPY

- ☐ Low blood counts / infection
- ☐ Sensory or motor neuropathy, potentially persistent
- ☐ Hypersensitivity / infusion reaction
- ☐ Hair loss and nail / skin changes
- ☐ Muscle and joint aches
- ☐ Fatigue and mouth / bowel symptoms
- ☐ Rare serious lung or cardiac effects

Confirm the indication, weekly schedule, hypersensitivity premedication, neuropathy assessment and any required off-label-use discussion.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00226 v9.

SACTCheck v0.83.0
Developed by Dr Paul O'Brien