NCCP 00216 · VERSION 8

Your treatment, explained

Colorectal and breast cancer · NCCP 00216 v8 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Capecitabine

21-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Stop capecitabine and contact oncology for significant diarrhoea, painful mouth or hand-foot symptoms affecting activity, vomiting that prevents drinking, fever or sudden illness. Do not restart after a toxicity interruption until the team advises.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp · hse

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- How the treatment fits together
- Your 21-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Tablets, visits and practical planning
- Your treatment aim and duration
- Taking capecitabine safely
- Tests and checks
- Recognise symptoms and act
- Daily life and other medicines
- Review, recovery and follow-up
- Supportive medicines
- Understanding the evidence
- Questions to ask
- Sources

UNDERSTAND YOUR TREATMENT


How the treatment works

Capecitabine is chemotherapy taken as tablets. NCCP 00216 covers several colorectal and breast cancer settings, including treatment after surgery and treatment of advanced disease. Ask which aim and duration apply to you.

ORAL CHEMOTHERAPY

Capecitabine

Xeloda



Becomes fluorouracil in the body

After you swallow capecitabine, your body changes it into fluorouracil. This interferes with the DNA and RNA processes cells use to grow. Normal cells can also be affected, which helps explain mouth soreness, bowel symptoms and changes in blood counts.

Swallowed tablet → Converted to fluorouracil → Cell growth disrupted

What to watch for

Stop tablets and contact the team for significant diarrhoea, painful mouth ulcers, hand-foot soreness affecting activities, fever or dehydration. Chest pain needs urgent assessment. Do not restart until instructed.

Sources: nccp · patient · smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Explore your existing anatomy and side-effect map

THE BIGGER TREATMENT PLAN

How the treatment fits together

ORAL CHEMOTHERAPY

21-day tablet cycle



capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

The treating team confirms the aim, duration and next steps. The visual pattern does not replace your prescription.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

Checks across the whole treatment course

The columns show the order of care, not equal lengths of time. Confirm who books each check, who reviews it and whom to call if it is missing.

Check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Blood count / kidneys / liver	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Tablets / hands and feet	Confirm written dose, tablet supply and hold instructions	Review adherence, mouth, bowel and skin symptoms	Days 1–14 tablets; Days 15–21 planned break unless team changes it	Do not restart held tablets or replace missed doses without advice	Return unused tablets as instructed; follow persistent symptoms
Heart checks	Review cardiac history; ECG when cardiac function is compromised or clinical findings warrant it. No automatic serial echocardiogram schedule is specified for this regimen.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00216 v8 · official regimen

THE WHOLE TREATMENT JOURNEY

Capecitabine: treatment and checks

Colorectal and breast cancer

capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before starting	Each treatment cycle	Between doses	Decision / response review	Completion / change
Treatment plan NCCP 00216	Confirm indication, medicines and duration	Give the agreed cycle only after review	Follow pump / tablet and symptom instructions	Continue, modify or stop after consultant review	Agree recovery and follow-up
Blood count / kidneys / liver NCCP 00216	Baseline bloods; DPD before fluoropyrimidine	Repeat before each cycle	Earlier tests if unwell, dehydrated or toxicity develops	Results and recovery guide any delay or dose change	Follow unresolved abnormalities
Symptoms and function NCCP 00216	Document existing symptoms and usual activity	Review worst symptoms since the last treatment	Use urgent contacts; do not wait for the next visit	Balance benefit, side effects and priorities	Agree recovery care and ongoing support
Tablets / hands and feet NCCP 00216	Confirm written dose, tablet supply and hold instructions	Review adherence, mouth, bowel and skin symptoms	Days 1–14 tablets; Days 15–21 planned break unless team changes it	Do not restart held tablets or replace missed doses without advice	Return unused tablets as instructed; follow persistent symptoms
Heart checks NCCP 00216	Review cardiac history; ECG when cardiac function is compromised or clinical findings warrant it. No automatic serial echocardiogram schedule is specified for this regimen.	Ask about new chest pain or breathlessness	Chest pain needs urgent assessment	Repeat heart tests if clinically indicated	Follow any heart problem; no fixed echo series
Response / scans NCCP 00216	Confirm cancer setting, disease assessment and intended duration	Review clinical benefit and symptoms	New symptoms may bring review forward	Consultant-directed scans and response assessment	Agree follow-up, completion or another treatment

What each check is for

1

Confirm the treatment aim

Before starting

Capecitabine is chemotherapy taken as tablets. NCCP 00216 covers several colorectal and breast cancer settings, including treatment after surgery and treatment of advanced disease. Ask which aim and duration apply to you.

Ask / act: Confirm the exact indication and intended duration.

Sources: NCCP 00216

2

Review before continuing

At each clinical decision point

Results, symptoms and benefit can change the plan.

Ask / act: Agree who reviews results and communicates the next step.

Sources: NCCP 00216

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?


This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-25 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.21

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.



Your written plan comes first: capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

Act when symptoms occur: Do not wait for the next visit or use the chart to decide a symptom is safe.

Symptoms can build: Bowel, mouth and hand-foot symptoms may accumulate and can continue during tablet-free days.

Across the course: The pattern can change between cycles. Dose changes and treatment breaks are made with the team.

Why these bands stay broad: Treatment bands show the NCCP schedule. Symptom bands are broad teaching illustrations: they do not represent measured daily risks or predict when you will feel well. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00216 v8 · official regimen ↗ · eviQ: taking capecitabine and managing symptoms (NCCP defines the Irish schedule) ↗

There are no guaranteed "safe days". Act on fever, sudden illness or a new serious symptom whenever it occurs. Do not wait for the next appointment.

Read the whole cycle as text

Day 1–14: Treatment days

capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription. Use the current tablet label. Swallow whole with water with or within 30 minutes of food; do not crush or cut tablets.

Day 15–21: Planned tablet-free days

Do not take leftover tablets to replace doses missed earlier. Side effects may continue during this interval. Keep the oncology contact plan available and confirm the next review before restarting.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.



STEP 1

Before the first treatment

Confirm the aim, likely course length, blood-test arrangements and a working urgent contact number. DPD testing is required before starting the fluoropyrimidine unless already completed.



STEP 2

Taking the tablets

Use the prescribed morning and evening dose within 30 minutes after food on the agreed treatment days. Do not make up a missed or vomited dose. Follow the team's hold instructions for significant diarrhoea, mouth soreness or painful hands and feet, and contact them promptly.



STEP 3

If treatment is delayed

Ask the unit to confirm the new treatment and blood-test dates. Re-anchor supportive medicines to the treatment actually given; do not take an extra or catch-up anticancer dose yourself.



STEP 4

Getting help between visits

Follow your oncology unit's urgent advice for fever, infection, severe diarrhoea, chest pain or difficulty breathing. Bring your treatment card and current medicines list.

Open practical treatment guide · Printable practical guide

Sources: NCCP 00216 v8 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Nausea prevention and rescue

The team agrees a rescue medicine if needed; routine daily anti-sickness prevention is not automatically required for this oral low-risk schedule.

2 Bowel symptoms

Report diarrhoea promptly and follow the written rescue plan. Severe symptoms, fever, dehydration or inability to drink need urgent advice.

3 Counts and recovery

White-cell growth-factor injections are prescribed according to individual risk and previous complications. They are not automatically required for every patient or every cycle.

4 Skin and mouth care

Use gentle skin and mouth care. Painful hands and feet or mouth sores can require a tablet hold and team review; do not continue simply to finish the pack.

Sources: nccp

Your treatment aim and duration

The same medicine can be used for different purposes.

Why this treatment?

Capecitabine is chemotherapy taken as tablets. NCCP 00216 covers several colorectal and breast cancer settings, including treatment after surgery and treatment of advanced disease. Ask which aim and duration apply to you.

Sources: nccp · patient

How long?

NCCP recommends six months / eight cycles for adjuvant colon or residual triple-negative breast cancer. For advanced disease, duration depends on response and tolerability.

Sources: nccp · patient

Alternatives and decisions

Discuss the expected benefit, burdens, reasonable alternatives and the option of supportive care. The best choice depends on your cancer, fitness and preferences.

Sources: nccp · patient

TREATMENT REFERENCE · 02

Taking capecitabine safely

Tablets are active chemotherapy, with a specific schedule.

Meals and doses

Use your current prescription. Swallow tablets whole with water with or within 30 minutes after food. Do not cut, chew or crush them; ask pharmacy if swallowing is difficult.

Sources: nccp · patient

Missed doses or vomiting

Do not replace a missed or vomited dose. Never double the next dose. Contact the team for repeated vomiting or uncertainty about the schedule.

Sources: nccp · patient

Pause and restart

Stop tablets for significant diarrhoea, painful mouth/hand-foot symptoms affecting activity, fever or dehydration and seek advice. A toxicity-related pause requires instructions before restart.

Sources: nccp · patient

Different tablet strengths

Different strengths may be combined to make your dose. A reduction changes tablet numbers: use the new label and do not calculate a change from this page.

Sources: nccp · patient

TREATMENT REFERENCE · 03

Tests and checks

Monitoring helps the team decide whether treatment can continue.

DPD testing

DPD helps break down fluoropyrimidine medicines. Low activity can cause severe toxicity; testing before treatment helps guide the plan but cannot prevent every serious reaction.

Sources: nccp · patient

Blood, kidney and liver results

Counts, kidney and liver function are checked before each cycle. Diarrhoea and poor intake can worsen kidney function and make tablets harder to tolerate.

Sources: nccp · patient

Your symptoms matter

Report bowel frequency, stoma output, mouth soreness, hand-foot pain, chest symptoms and what daily activities you can no longer manage.

Sources: nccp · patient

Additional checks

The team sets the scan or follow-up plan for your particular treatment aim.

Sources: nccp · patient

TREATMENT REFERENCE · 04

Recognise symptoms and act

Do not wait for the next appointment if you feel unwell.

Bowel and hydration

Report diarrhoea or increased stoma output early. Inability to drink, dizziness, very little urine, severe pain or blood in stool needs prompt assessment.

Sources: nccp · patient

Hands, feet and mouth

Pain, cracking, peeling or ulcers that affect normal activities or eating/drinking need advice. Gentle preventive care helps, but does not replace reporting.

Sources: nccp · patient

Chest symptoms and infection

Stop capecitabine and seek urgent assessment for chest pain. Fever, shaking chills or sudden illness needs immediate oncology contact, even if the temperature is normal.

Sources: nccp · patient

Treatment-specific effects

Severe toxicity can happen early. Do not assume that tablets are mild treatment or that a normal DPD result makes symptoms safe to ignore.

Sources: nccp · patient

TREATMENT REFERENCE · 05

Daily life and other medicines

Make the plan fit your practical needs.

Eating, drinking and activity

Aim for tolerable meals and fluids; contact the team if intake falls. Gentle activity and rest can be adjusted to your energy. Ask for dietetic help if needed.

Sources: nccp · patient

Medicine and supplement checks

Check new medicines with oncology pharmacy. Warfarin can require closer INR checks; phenytoin levels may need monitoring. Brivudine is an important prohibited interaction.

Sources: nccp · patient

Family, work and fertility

Ask about infection exposure, work, intimacy, contraception, fertility and any caring responsibilities. Use the medicine-specific duration of precautions given by your team.

Sources: nccp · patient

Procedures and travel

Plan transport and support around appointments. Ask which symptoms or blood-count problems affect travel and when to contact the service.

Sources: nccp · patient

TREATMENT REFERENCE · 06

Review, recovery and follow-up

Your plan can change as treatment progresses.

Dose changes are individual

A pause or reduction may be needed because of symptoms, blood results or kidney function. The team considers the treatment aim and other options.

Sources: nccp · patient

After the last dose

Keep reporting new or persistent symptoms after treatment ends. Your team confirms follow-up, scans and recovery support; the timeline is not a forecast.

Sources: nccp · patient

Prepare for review

Bring questions, symptoms and missed-dose records. Tell the team what matters most to you and ask them to explain the next decision.

Sources: nccp · patient

Understanding the evidence

Ask which evidence matches your cancer, treatment aim and medicines.

In stage III colon cancer after surgery, capecitabine was at least as effective as the intravenous fluorouracil schedule studied. This comparison does not measure benefit against no treatment, and does not estimate benefit for breast cancer or cancer that has spread.

Sources: trial

PREPARE FOR THE CONVERSATION

Questions worth asking

What is the aim of my treatment?

Capecitabine is chemotherapy taken as tablets. NCCP 00216 covers several colorectal and breast cancer settings, including treatment after surgery and treatment of advanced disease. Ask which aim and duration apply to you.

Which days do I take the tablets?

capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

What if I forget a dose or vomit?

Do not take an extra dose. Take the next planned dose at its usual time unless treatment has been stopped; report repeated vomiting and check with the team if unsure.

When should I stop tablets and call?

Stop capecitabine and contact oncology for significant diarrhoea, painful mouth or hand-foot symptoms affecting activity, vomiting that prevents drinking, fever or sudden illness. Do not restart after a toxicity interruption until the team advises.

Why do I need blood tests if I take tablets at home?

Blood counts, kidney and liver results, together with symptoms, guide whether to continue and at what dose.

Do I need a port or a pump?

No take-home chemotherapy pump is used. Capecitabine is swallowed, so a central line is not required just to give it.

What happens after treatment?

Ask for your review and follow-up plan. Continue reporting symptoms after the last dose; further therapy depends on the treatment aim and response.

> **Try explaining the plan back:** Which days do you take tablets? What do you do if you miss a dose? Which symptoms mean stop tablets and call? Who do you contact out of hours?

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00216 v8. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00216 v8 · official regimen
- eviQ: taking capecitabine and managing symptoms (NCCP defines the Irish schedule)
- Capecitabine product information: medicines, interactions and precautions
- HSE: chemotherapy side effects and urgent advice
- X-ACT: primary trial report

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-10-25.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6558/216_v8_Capecitabine.pdf

https://www.eviq.org.au/medical-oncology/colorectal/adjuvant-and-neoadjuvant/74-colorectal-adjuvant-capecitabine/patient-information

https://www.medicines.org.uk/emc/product/14545/smpc

https://www2.hse.ie/conditions/chemotherapy/side-effects/

https://pubmed.ncbi.nlm.nih.gov/21896539/