SACTCHECK · NCCP 00203

My complete treatment handbook

Docetaxel alone for previously treated non-small-cell lung cancer

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

Contents

1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 21
4. Supportive medicines and care — page 24
5. Visits, access and practical arrangements — page 25
6. Your treatment passport and symptom record — page 26
7. Wallet card and phone instructions — page 33

SACTCheck v0.84.33 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

Docetaxel alone for previously treated non-small-cell lung cancer

To help control previously treated non-small-cell lung cancer and relieve symptoms.

Lung · previously treated advanced NSCLC · NCCP 00203 v7 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Docetaxel by drip on Day 1, repeated every 21 days, for up to six cycles. Treatment may stop earlier if it is not helping or side effects are unacceptable. Your team confirms the prescribed course.

Before starting	Treatment every 21 days	After treatment
Confirm your blood tests, current medicines and steroid plan.	Day 1 drip, then time at home. Up to six cycles, with a review before each.	Review benefit, recovery and the next care plan with your team.

A typical 21-day cycle

Day 1	Day 2	Day 3–20	Day 21
Treatment day	Day after treatment	Between visits	Prepare for the next cycle

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before starting	**Blood count, kidneys and liver** Required by regimen	Before the first dose
During treatment	**Blood count, kidneys and liver** Required by regimen	Before every 21-day cycle
During / after	**Tumour markers, scans or extra tests** If clinically indicated	Scans: team-agreed timing. Markers: only if relevant; no routine schedule.
After treatment	**Recovery and follow-up** Team-agreed plan	Team agrees timing; follow ongoing symptoms or abnormal results

Between treatments

- Day 2: finish prescribed steroid. Between visits, watch for infection, bowel changes, numbness or swelling; timing varies.
- Before another cycle: blood tests and symptom review. Delays or early stopping may be needed.

Take the prescribed steroid from the day before each infusion. Tell the team if any doses were missed. Bring your medicine list and report symptoms since the last visit. Ask when your next blood tests and cancer review will be.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hands / feet

Docetaxel

Report numbness, weakness or difficulty walking or using your hands.

3 Hair / scalp

Docetaxel

Hair loss can occur; ask about support before treatment.

5 Mouth


Docetaxel

Report painful ulcers, difficulty drinking or poor intake.

7 Skin / swelling

Docetaxel

Report swelling; blistering or peeling skin needs urgent advice.



2 Blood / infection

Docetaxel

Fever, shivers or sudden illness need immediate contact.

4 Breathing

Docetaxel

New or worsening cough or breathlessness needs urgent assessment.

6 Tummy / bowel

Docetaxel

Severe tummy pain or rapidly worsening diarrhoea needs urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call the oncology contact number immediately for fever, shivers or sudden illness, severe or new breathing difficulty, chest pain, fainting, severe abdominal pain or persistent diarrhoea. Tell staff immediately about infusion symptoms. Report new numbness, weakness, painful skin changes, swelling or visual changes promptly; do not wait for the next cycle. Severe tummy pain, rapidly worsening diarrhoea, blistering or peeling skin, or new breathing symptoms also need urgent assessment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

Lung · previously treated advanced NSCLC · NCCP 00203 v7 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card SACTCheck

Name: _____

Docetaxel
Day 1 every 21 days; up to 6 cycles
Phase: _____ Planned cycles: 6
Daytime (EXAMPLE) 087 12345678
Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1	C2	C3
Date / initials	Date / initials	Date / initials
C4	C5	C6
Date / initials	Date / initials	Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

NCCP 00203 · VERSION 7

Your treatment, explained

Lung · previously treated advanced NSCLC · NCCP 00203 v7 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Docetaxel alone for previously treated non-small-cell lung cancer

21-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Call the oncology contact number immediately for fever, shivers or sudden illness, severe or new breathing difficulty, chest pain, fainting, severe abdominal pain or persistent diarrhoea. Tell staff immediately about infusion symptoms. Report new numbness, weakness, painful skin changes, swelling or visual changes promptly; do not wait for the next cycle. Severe tummy pain, rapidly worsening diarrhoea, blistering or peeling skin, or new breathing symptoms also need urgent assessment.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp · smpc

Choose a topic to read

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- Treatment pattern
- Your 21-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Tablets, visits and practical planning
- Your lung treatment plan
- Checks through treatment
- Practical support
- Supportive medicines
- Questions to ask
- Sources

UNDERSTAND YOUR TREATMENT

How the medicines work

Treatment aims to slow or shrink lung cancer and relieve symptoms after previous chemotherapy. Expected benefit, side effects, other options and your priorities guide the decision.

TAXANE CHEMOTHERAPY

Docetaxel



Disrupts the cell's division machinery

Docetaxel stabilises tiny structures called microtubules. Cells need these to separate their chromosomes during division; disrupting the process can stop cancer cells multiplying. Healthy dividing cells are also affected.

Microtubules stabilised → Division disrupted → Cell growth affected

What to watch for

Take prescribed steroid premedication. Report fever, lasting numbness, swelling, sore nails or infusion symptoms.

Sources: nccp · smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Explore your existing anatomy and side-effect map

THE BIGGER TREATMENT PLAN

Treatment pattern

EVERY 21 DAYS

Docetaxel monotherapy

Cycle 1 · Cycle 2 · Cycle 3 · Cycle 4 · Cycle 5 · Cycle 6

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.

The team confirms your actual prescription and appointments. This map explains the pattern and does not decide whether a dose can be given.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Relative stages explain the order of care; they are not equally spaced appointments. FBC, kidney and liver profile at baseline and before every 21-day cycle; review symptoms, neuropathy, fluid retention and the steroid plan. Confirm remaining cycles against the six-cycle NSCLC maximum.

Check	Before starting	Early treatment	Ongoing treatment	Pause / dose change	Completion / next decision
Blood / kidney / liver	Baseline FBC, kidney and liver tests	Before each 21-day cycle	Repeat before each planned dose	Check recovery before restarting	Follow unresolved abnormalities
Symptoms / premedication	Review symptoms, medicines and reaction history	Check steroid doses, swelling and neuropathy	Review before every treatment	Record worst and current symptoms; clinician confirms changes	Review persistent symptoms and recovery
Heart / breathing	Assess history and symptoms	Infusion observations and symptom review	Extra tests if symptoms or risk require	Urgent assessment for chest pain, fainting or new breathlessness	Agree follow-up for unresolved symptoms
Cancer / course	Confirm 00203a NSCLC after prior chemotherapy	Agree response review and actual appointment plan	Up to six cycles; stop earlier if needed	Reassess benefit, toxicity and priorities	Agree next treatment or supportive-care plan

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00203 v7 · official regimen · EMA Taxotere product information; retrieved 2 November 2026

THE WHOLE TREATMENT JOURNEY

Docetaxel alone for previously treated non-small-cell lung cancer: treatment and checks

Lung · previously treated advanced NSCLC

Docetaxel by drip on Day 1, repeated every 21 days, for up to six cycles. Treatment may stop earlier if it is not helping or side effects are unacceptable. Your team confirms the prescribed course.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before starting	Early treatment	Ongoing treatment	Pause / dose change	Completion / next decision
Treatment plan NCCP 00203	Confirm indication and prescription	Day 1 every 21 days	Maximum six cycles for NSCLC	Clinician review before restart	Agree the next plan
Blood / kidney / liver NCCP 00203 · SMPC	Baseline FBC, kidney and liver tests	Before each 21-day cycle	Repeat before each planned dose	Check recovery before restarting	Follow unresolved abnormalities
Symptoms / premedication NCCP 00203 · SMPC	Review symptoms, medicines and reaction history	Check steroid doses, swelling and neuropathy	Review before every treatment	Record worst and current symptoms; clinician confirms changes	Review persistent symptoms and recovery
Heart / breathing NCCP 00203 · SMPC	Assess history and symptoms	Infusion observations and symptom review	Extra tests if symptoms or risk require	Urgent assessment for chest pain, fainting or new breathlessness	Agree follow-up for unresolved symptoms
Cancer / course NCCP 00203 · SMPC	Confirm 00203a NSCLC after prior chemotherapy	Agree response review and actual appointment plan	Up to six cycles; stop earlier if needed	Reassess benefit, toxicity and priorities	Agree next treatment or supportive-care plan

What each check is for

1

Safety review

Before each 21-day cycle

FBC, kidney and liver profile at baseline and before every 21-day cycle; review symptoms, neuropathy, fluid retention and the steroid plan. Confirm remaining cycles against the six-cycle NSCLC maximum.

Ask / act: The team reviews results, current symptoms and the prescription.

Sources: NCCP 00203 · SMPC

2

Benefit and treatment choice

At consultant-agreed response reviews

A historical randomised trial found that docetaxel 75 mg/m² every three weeks prolonged survival compared with supportive care alone in selected people whose NSCLC had progressed after platinum chemotherapy. Modern previous treatment and available alternatives differ, so the trial cannot predict an individual outcome.

Ask / act: Discuss continuing, changing or stopping treatment according to benefit, toxicity and priorities.

Sources: NCCP 00203 · TRIAL

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.29

One 21-day cycle

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.

Day of cycle	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Chemotherapy · Day 1	IV																				
Administration reaction	Report during or after any visit																				
Nausea / appetite	May occur early or later																				
Blood counts / infection	Risk varies; fever needs prompt advice																				
Fatigue / mouth changes	Variable timing and severity																				
Nerves / skin / nails	Can build across doses																				

Six-cycle limit: The NSCLC source plan has at most six cycles. Progression or unacceptable side effects may stop treatment sooner.

Between visits: Symptoms may occur on any day. Do not wait for the next infusion to report urgent problems.

No dates to enter: This diagram explains the sequence; your team books actual visits and tests.

Why these bands stay broad: These broad symptom bands are educational possibilities, not measured risk curves. Infusion days are the planned schedule; delays require a new clinical plan. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00203 v7 · official regimen ↗

There are no guaranteed "safe days". Fever, shivers, sudden illness, severe breathing changes or chest pain need prompt assessment on any day.

Read the whole cycle as text

Day 1: Treatment day

Your blood results, symptoms and steroid preparation are checked before docetaxel. The infusion takes about one hour; the full visit is longer. Tell the nurse immediately about discomfort at the drip, flushing, breathing symptoms or feeling faint.

Day 2: Day after treatment

Finish the prescribed three-day steroid course and use anti-sickness medicine as directed. Ask the team if a steroid dose was missed; do not invent a replacement dose.

Day 3–20: Between visits

Watch for infection, mouth soreness, fatigue, bowel changes, numbness or swelling. Symptoms and recovery vary; no day is guaranteed to be risk-free. Contact the team promptly about worrying symptoms, even if your temperature is normal.

Day 21: Prepare for the next cycle

The team arranges blood tests and reviews recovery and benefit before another dose. Your next treatment day depends on this review. Check the next appointment and the steroid start date. Do not use a passport stamp as permission for another dose.

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.



STEP 1

Before the visit

Check the oral steroid schedule, blood-test appointment and transport. Contact the unit before attending if premedication was missed or you are unwell.



STEP 2

At the unit

Docetaxel is given through a cannula or suitable existing IV line over about one hour. The appointment also includes checks, preparation, premedication and observation and is longer than the infusion itself. This regimen has no routine take-home pump.



STEP 3

At home

Keep the contact number, medicine list and symptom record available. Ask who will arrange your next blood test, scan and review. Sedating medicines may affect driving.

Planned course

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.

Open practical treatment guide · Printable practical guide

Sources: NCCP 00203 v7 · official regimen · EMA Taxotere product information; retrieved 2 November 2026. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Premedication

Take dexamethasone exactly as prescribed before and after docetaxel. Tell the unit about missed doses before infusion; any alternative needs the prescriber’s decision. Do not double a dose or add another steroid course yourself.

2 Nausea, bowel symptoms and pain

Use your individual anti-sickness and pain plan. Contact the team if symptoms are uncontrolled or you cannot drink. Antihistamines and other support medicines may cause drowsiness; arrange transport if advised.

3 Nerves and daily activities

Report tingling, numbness or weakness early, including effects on buttons, writing, walking or balance. Protect hands and feet and take care with hot surfaces if sensation changes.

4 Infection and blood counts

Each treatment needs its own blood and symptom review. A growth-factor injection may be prescribed after individual risk assessment; follow its separate instructions. Fever or sudden illness needs immediate contact on any day.

Sources: nccp · smpc · antiemetic

TREATMENT REFERENCE · 01

Your lung treatment plan

Treatment aims to slow or shrink lung cancer and relieve symptoms after previous chemotherapy. Expected benefit, side effects, other options and your priorities guide the decision.

Exact cancer setting

NCCP 00203a: locally advanced or metastatic non-small-cell lung cancer after failure of prior chemotherapy. This is docetaxel alone; a platinum or other partner needs its own regimen.

Sources: nccp

Course and visits

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.

Sources: nccp

What the evidence can tell us

A historical randomised trial found that docetaxel 75 mg/m² every three weeks prolonged survival compared with supportive care alone in selected people whose NSCLC had progressed after platinum chemotherapy. Modern previous treatment and available alternatives differ, so the trial cannot predict an individual outcome.

Sources: trial

TREATMENT REFERENCE · 02

Checks through treatment

Your team arranges tests, reviews results and confirms each dose.

Blood tests and symptoms

FBC, kidney and liver profile at baseline and before every 21-day cycle; review symptoms, neuropathy, fluid retention and the steroid plan. Confirm remaining cycles against the six-cycle NSCLC maximum.

Sources: nccp

Heart and breathing

Assess cardiac history and symptoms; this monotherapy source does not prescribe routine serial echocardiograms. New breathlessness, chest pain or fainting needs assessment.

Sources: nccp · smpc

Response review

The consultant agrees when scans and other disease checks are due. New or worsening symptoms may need assessment sooner. There is no automatic fixed scan interval in this regimen.

Sources: nccp

Practical support

Bring your medicine list and symptom record to every review.

Medicines and vaccines

Check new prescriptions, over-the-counter medicines, supplements and vaccine timing with the oncology pharmacist. Interactions and changes in organ function can alter the treatment plan.

Sources: smpc

Fertility and contraception

Discuss fertility preservation before treatment where relevant. Ask for written contraception and breastfeeding advice for your actual product and the period after treatment. Tell the team promptly about a possible pregnancy.

Sources: smpc

Support and priorities

Ask about symptom control, nutrition, rehabilitation, emotional support and help at home. Review how the treatment burden and possible benefit fit your priorities.

Sources: nccp · trial

Why this treatment may be chosen

A historical randomised trial found that docetaxel 75 mg/m^2 every three weeks prolonged survival compared with supportive care alone in selected people whose NSCLC had progressed after platinum chemotherapy. Modern previous treatment and available alternatives differ, so the trial cannot predict an individual outcome.

Discuss possible benefit, alternatives and uncertainty for your own situation. The treatment aim is cancer control and symptom relief.

Sources: nccp · trial

PREPARE FOR THE CONVERSATION

Questions worth asking

What benefit is realistic for me and what other options are available?

Treatment aims to slow or shrink lung cancer and relieve symptoms after previous chemotherapy. Expected benefit, side effects, other options and your priorities guide the decision. A historical randomised trial found that docetaxel 75 mg/m² every three weeks prolonged survival compared with supportive care alone in selected people whose NSCLC had progressed after platinum chemotherapy. Modern previous treatment and available alternatives differ, so the trial cannot predict an individual outcome.

What are my confirmed infusion days, dose and planned review points?

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.

Who arranges and reviews my blood tests and scans?

FBC, kidney and liver profile at baseline and before every 21-day cycle; review symptoms, neuropathy, fluid retention and the steroid plan. Confirm remaining cycles against the six-cycle NSCLC maximum. Ask the team who books each test and who checks the result.

What premedication and anti-sickness plan should I follow?

Follow the written plan for your exact prescription. Report missed premedication before infusion; confirm prevention and rescue medicines with the pharmacist.

Which symptoms need an immediate call, and which number should I use?

Call the oncology contact number immediately for fever, shivers or sudden illness, severe or new breathing difficulty, chest pain, fainting, severe abdominal pain or persistent diarrhoea. Tell staff immediately about infusion symptoms. Report new numbness, weakness, painful skin changes, swelling or visual changes promptly; do not wait for the next cycle. Keep the unit's daytime and out-of-hours numbers on your treatment card.

> **Try explaining the plan back:** Can you explain the treatment aim, your next visit, the blood-test plan, premedication and when to call for urgent help?

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00203 v7. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00203 v7 · official regimen
- EMA Taxotere product information; retrieved 2 November 2026
- Shepherd et al., TAX317, JCO 2000; previously treated NSCLC
- NCCP NCIS medical-oncology antiemetics V8
- Fossella et al., TAX320, JCO 2000

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-11-02.

This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6543/203_DOCEtaxel_75-21day.pdf

https://www.ema.europa.eu/en/documents/product-information/taxotere-epar-product-information_en.pdf

https://pubmed.ncbi.nlm.nih.gov/10811675/

https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf

https://pubmed.ncbi.nlm.nih.gov/10856094/

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Relative stages explain the order of care; they are not equally spaced appointments. FBC, kidney and liver profile at baseline and before every 21-day cycle; review symptoms, neuropathy, fluid retention and the steroid plan. Confirm remaining cycles against the six-cycle NSCLC maximum.

Check	Before starting	Early treatment	Ongoing treatment	Pause / dose change	Completion / next decision
Blood / kidney / liver	Baseline FBC, kidney and liver tests	Before each 21-day cycle	Repeat before each planned dose	Check recovery before restarting	Follow unresolved abnormalities
Symptoms / premedication	Review symptoms, medicines and reaction history	Check steroid doses, swelling and neuropathy	Review before every treatment	Record worst and current symptoms; clinician confirms changes	Review persistent symptoms and recovery
Heart / breathing	Assess history and symptoms	Infusion observations and symptom review	Extra tests if symptoms or risk require	Urgent assessment for chest pain, fainting or new breathlessness	Agree follow-up for unresolved symptoms
Cancer / course	Confirm 00203a NSCLC after prior chemotherapy	Agree response review and actual appointment plan	Up to six cycles; stop earlier if needed	Reassess benefit, toxicity and priorities	Agree next treatment or supportive-care plan

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00203 v7 · official regimen · EMA Taxotere product information; retrieved 2 November 2026

THE WHOLE TREATMENT JOURNEY

Docetaxel alone for previously treated non-small-cell lung cancer: treatment and checks

Lung · previously treated advanced NSCLC

Docetaxel by drip on Day 1, repeated every 21 days, for up to six cycles. Treatment may stop earlier if it is not helping or side effects are unacceptable. Your team confirms the prescribed course.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Early treatment	Ongoing treatment	Pause / dose change	Completion / next decision
Treatment plan NCCP 00203	Confirm indication and prescription	Day 1 every 21 days	Maximum six cycles for NSCLC	Clinician review before restart	Agree the next plan
Blood / kidney / liver NCCP 00203 · SMPC	Baseline FBC, kidney and liver tests	Before each 21-day cycle	Repeat before each planned dose	Check recovery before restarting	Follow unresolved abnormalities
Symptoms / premedication NCCP 00203 · SMPC	Review symptoms, medicines and reaction history	Check steroid doses, swelling and neuropathy	Review before every treatment	Record worst and current symptoms; clinician confirms changes	Review persistent symptoms and recovery
Heart / breathing NCCP 00203 · SMPC	Assess history and symptoms	Infusion observations and symptom review	Extra tests if symptoms or risk require	Urgent assessment for chest pain, fainting or new breathlessness	Agree follow-up for unresolved symptoms
Cancer / course NCCP 00203 · SMPC	Confirm 00203a NSCLC after prior chemotherapy	Agree response review and actual appointment plan	Up to six cycles; stop earlier if needed	Reassess benefit, toxicity and priorities	Agree next treatment or supportive-care plan

What each check is for

1 Safety review

Before each 21-day cycle

FBC, kidney and liver profile at baseline and before every 21-day cycle; review symptoms, neuropathy, fluid retention and the steroid plan. Confirm remaining cycles against the six-cycle NSCLC maximum.

Ask / act: The team reviews results, current symptoms and the prescription.

Sources: NCCP 00203 · SMPC

2 Benefit and treatment choice

At consultant-agreed response reviews

A historical randomised trial found that docetaxel 75 mg/m² every three weeks prolonged survival compared with supportive care alone in selected people whose NSCLC had progressed after platinum chemotherapy. Modern previous treatment and available alternatives differ, so the trial cannot predict an individual outcome.

Ask / act: Discuss continuing, changing or stopping treatment according to benefit, toxicity and priorities.

Sources: NCCP 00203 · TRIAL

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.29

Docetaxel alone for previously treated non-small-cell lung cancer

Treatment scope · NCCP 00203 NCCP 00203a: locally advanced or metastatic non-small-cell lung cancer after failure of prior chemotherapy. This is docetaxel alone; a platinum or other partner needs its own regimen.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Docetaxel · lung supportive care** — low
  Docetaxel

https://sactcheck.com/patient/00203/lung/supportive-care.html

Docetaxel alone for previously treated non-small-cell lung cancer

NCCP 00203 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.

STEP 1

Before the visit

Check the oral steroid schedule, blood-test appointment and transport. Contact the unit before attending if premedication was missed or you are unwell.

STEP 2

At the unit

Docetaxel is given through a cannula or suitable existing IV line over about one hour. The appointment also includes checks, preparation, premedication and observation and is longer than the infusion itself. This regimen has no routine take-home pump.

STEP 3

At home

Keep the contact number, medicine list and symptom record available. Ask who will arrange your next blood test, scan and review. Sedating medicines may affect driving.

Planned course

Docetaxel 75 mg/m² is infused on Day 1 every 21 days. For this NSCLC indication, the NCCP plan has a maximum of six cycles; treatment stops earlier for progression or unacceptable side effects. Your team confirms your prescribed dose and course.

Sources: NCCP 00203 v7 · official regimen · EMA Taxotere product information; retrieved 2 November 2026. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

Docetaxel alone for previously treated non-small-cell lung cancer

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Docetaxel by drip on Day 1, repeated every 21 days, for up to six cycles. Treatment may stop earlier if it is not helping or side effects are unacceptable. Your team confirms the prescribed course.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital’s instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00203/lung/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Docetaxel alone for previously treated non-small-cell lung cancer

Cycle number: _____ Start date: _____ Next review: _____

Docetaxel · cycle 1 of up to six · copy this sheet for another cycle of the same phase.

Docetaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Docetaxel alone for previously treated non-small-cell lung cancer

Cycle number: _____ Start date: _____ Next review: _____

Docetaxel · cycle 2 of up to six · copy this sheet for another cycle of the same phase.

Docetaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Docetaxel alone for previously treated non-small-cell lung cancer

Cycle number: _____ Start date: _____ Next review: _____

Docetaxel · cycle 3 of up to six · copy this sheet for another cycle of the same phase.

Docetaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Docetaxel alone for previously treated non-small-cell lung cancer

Cycle number: _____ Start date: _____ Next review: _____

Docetaxel · cycle 4 of up to six · copy this sheet for another cycle of the same phase.

Docetaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Docetaxel alone for previously treated non-small-cell lung cancer

Cycle number: _____ Start date: _____ Next review: _____

Docetaxel · cycle 5 of up to six · copy this sheet for another cycle of the same phase.

Docetaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Docetaxel alone for previously treated non-small-cell lung cancer

Cycle number: _____ Start date: _____ Next review: _____

Docetaxel · cycle 6 of up to six · copy this sheet for another cycle of the same phase.

Docetaxel · Taxane chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

Lung · previously treated advanced NSCLC · NCCP 00203 v7 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card SACTCheck

Name: _____

Docetaxel

Day 1 every 21 days; up to 6 cycles

Phase: _____ Planned cycles: 6

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1 Date / initials

C2 Date / initials

C3 Date / initials

C4 Date / initials

C5 Date / initials

C6 Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.32 · NCCP 00203 v7 · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.