SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Dose-dense AC → paclitaxel every 2 weeks

neoadjuvant or adjuvant high-risk breast cancer · NCCP 00278 · version 7 · 14-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Part of an individual breast cancer plan to reduce the risk of recurrence, and to shrink the cancer before surgery where that is the agreed setting.

Schedule
Four AC cycles 14 days apart, followed by four paclitaxel doses 14 days apart. Growth-factor support accompanies the dose-dense course.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Required blood tests and cardiac monitoring discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Doxorubicin
ANTHRACYCLINE CHEMOTHERAPY

- ☐ Low counts, infection and mouth ulcers
- ☐ Nausea, hair loss and fatigue
- ☐ Heart damage, sometimes delayed
- ☐ Severe tissue injury if the infusion leaks
- ☐ Rare later blood cancers

Cyclophosphamide
ALKYLATING CHEMOTHERAPY

- ☐ Low counts and infection
- ☐ Nausea, vomiting and hair loss
- ☐ Bladder irritation / bleeding
- ☐ Fertility effects and menopause
- ☐ Rare later blood cancers

Paclitaxel
TAXANE CHEMOTHERAPY

- ☐ Low blood counts and infection
- ☐ Numbness / tingling, sometimes lasting
- ☐ Hair loss, muscle or joint pain
- ☐ Infusion reaction
- ☐ Mouth, bowel, skin and nail effects

Use this information to discuss the treatment goal, alternatives, phase changes, important risks and monitoring with your team. It supports the approved local consent process.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00278 v7.

SACTCheck v0.84.17
Developed by Dr Paul O'Brien