SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Pemetrexed + Carboplatin

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Pleural mesothelioma / non-squamous NSCLC · chemotherapy component · NCCP 00318 · version 6 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Pemetrexed + Carboplatin is chemotherapy used for selected people with unresectable pleural mesothelioma or non-squamous non-small-cell lung cancer. The aim may be disease control or, within a separate combined preoperative plan, support for surgery. Ask your team to explain the expected benefit for your own diagnosis.

Schedule
Both chemotherapy medicines are planned on Day 1 of a 21-day cycle. For pleural mesothelioma, the source describes up to 4–6 cycles. For advanced non-squamous NSCLC, it describes 4–6 combination cycles followed, when appropriate, by a separate pemetrexed maintenance plan (NCCP 00222). The selected preoperative nivolumab combination uses three cycles and a separate immunotherapy plan (NCCP 00849). Your team confirms which course applies and may stop or change it earlier.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Pemetrexed
ANTIFOLATE CHEMOTHERAPY

- ☐ Low blood counts, infection and bleeding
- ☐ Nausea, vomiting and fatigue
- ☐ Mouth soreness and diarrhoea
- ☐ Kidney problems
- ☐ Allergic or infusion reactions
- ☐ Skin rash; rarely serious skin reactions
- ☐ Rare serious lung inflammation

Carboplatin
PLATINUM CHEMOTHERAPY

- ☐ Low blood counts, infection and bleeding
- ☐ Nausea, vomiting and fatigue
- ☐ Mouth soreness and diarrhoea
- ☐ Kidney problems
- ☐ Allergic or infusion reactions
- ☐ Hearing changes or tinnitus, which can persist
- ☐ Numbness or tingling, which can persist

Confirm the treatment aim and indication, likely benefit and uncertainty, reasonable alternatives including not proceeding, serious and lasting risks, kidney/hearing/nerve problems, fertility and pregnancy advice, vitamin and steroid plans, and the chance of delays or dose changes. For mesothelioma and advanced NSCLC, the source identifies this carboplatin combination as off-label in Ireland; ask the team to explain this and document consent under hospital policy.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00318 v6.

SACTCheck v0.84.28
Developed by Dr Paul O'Brien