Carboplatin and paclitaxel · lung cancer

After surgery, treatment aims to reduce the chance of lung cancer returning. For advanced lung cancer, it aims to control cancer and help symptoms. Your team confirms which aim applies to you.

Your treatment plan

Both medicines on Day 1, every 21 days (three weeks), for four planned cycles.

1. **Before starting**
Confirm why treatment is being offered, blood tests, medicines and whom to call.

2. **Four treatment cycles**
One treatment visit every three weeks, with time at home between visits. The team may adjust or delay doses.

3. **After the course**
Review recovery and agree the next steps. Other treatment or scans follow your own plan.

Before and between visits

- Bring your medicine list and report new symptoms.
- Confirm blood tests and your allergy-prevention medicines.
- Keep the daytime and out-of-hours numbers with you.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately for temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions.

Cannot reach the team? Seek urgent medical assessment. Do not wait for the next appointment.

Your contact plan

Ask your team to complete these before you leave.

Hospital / team _____

Daytime number and hours _____

Evening / weekend number _____

If unavailable, where to go _____

Keep this with your consent information. Ask your team about anything unclear.

SACTCheck · v0.84.28 · Source: NCCP 00304 v8. Feasibility build: independent clinical and patient review pending. Your team confirms your personal plan.

What to watch for

Use the picture and symptom descriptions together. You do not need to know which medicine caused a symptom before calling.

1. Hair and skin

Paclitaxel

Hair loss and skin or nail changes can occur. Report severe rash or blistering urgently.

2. Mouth

Both medicines

Report painful sores or difficulty eating or drinking.

3. Blood and infection

Both medicines

Fever or sudden illness needs an immediate call, even between visits.

4. Stomach and bowels

Both medicines

Use prescribed anti-sickness medicines; call for severe pain, persistent vomiting or diarrhoea.

5. Kidneys and urine

Carboplatin

Blood tests guide the dose; report reduced urine or poor drinking.

6. Nerves and hearing


Both medicines

Report tingling, numbness, weakness, balance or hearing changes early.

7. Liver

Paclitaxel

Blood tests monitor liver function; report yellow eyes or dark urine.



Selected effects, not every possible effect or a prediction. Match each number to its symptom box. Blood, skin and nerve markers are schematic.

Call your oncology team immediately for fever, shivering, sudden illness, new breathing difficulty, chest pain, unusual bleeding, severe abdominal pain or inability to drink. Report new numbness, weakness or hearing change promptly. Follow the unit’s emergency instructions and do not wait for the next visit.

My SACT card

Carboplatin and paclitaxel · lung cancer
Day 1 every 21 days; four planned cycles.

Receiving carboplatin and paclitaxel. Fever or sudden illness needs urgent oncology advice.

Name: _____
Hospital: _____
Oncology phone: _____
Out of hours: _____

Show this card when seeking care. It does not replace a medicine-specific alert card.

More information, when you want it

Scan for medicines, tests, daily life and your treatment record.

https://sactcheck.com/patient/00304/

SACTCheck · v0.84.28 · Source: NCCP 00304 v8. Feasibility build: independent clinical and patient review pending. Your team confirms your personal plan.