Gemcitabine and cisplatin · lung cancer

NCCP 00281 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before the visit

Confirm blood tests, transport, medicine instructions and contact numbers.

STEP 2

Time in the unit

Gemcitabine runs through a drip over 30 minutes on Days 1 and 8. Cisplatin runs over 60 minutes on Day 1, with fluids and electrolytes before and afterwards according to your prescribed plan. Blood tests, pharmacy preparation, support medicines and review add to the visit time.

STEP 3

Access

Suitable IV access is needed. This plan does not use a routine take-home infusion pump.

STEP 4

Home support

Follow one written anti-sickness prescription, including any tablets for home and rescue treatment. Day 8 gemcitabine has a separate lower-risk anti-sickness plan. Cisplatin needs prescribed IV hydration, electrolyte replacement and kidney, hearing and nerve checks. Follow your own fluid advice; tell the team if you have heart failure or a fluid restriction.

STEP 5

Next appointment

Confirm the next Day 1 or Day 8 visit and any response scan before leaving.

Sources: NCCP 00281 v7 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.