NCCP 00310 · clinician draft

Gemcitabine and carboplatin · lung cancer

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Day 8 · gemcitabine alone

Active medicines: gemcitabine

Emetogenic risk: low

Medicine	Dose / route	When
Dexamethasone	8 mg orally	Day 8 before gemcitabine if prescribed NCIS low-risk reference; individualise and reconcile with any other steroid course.

Additional phase-specific support

- Follow one written anti-sickness prescription, including any tablets for home and rescue treatment. Day 8 gemcitabine has a separate lower-risk anti-sickness plan.Carboplatin needs a reliable kidney-function result for dosing. Report unusual bruising or bleeding promptly.
- Follow one reconciled prescription; the clinical support sheet is a reference, not an instruction to self-prescribe.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00310 v8 · pNCCP 00310 supportive care; NCIS V 8 Table 1 · NCCP NCIS antiemetics V 8

Sources checked 2026-11-02 · SACTCheck v0.84.28 · Regimen-level drafts; prescriber and oncology-pharmacy verification required