Vinorelbine and cisplatin · 21-day cycle

After surgery, treatment aims to lower the chance of cancer returning. For advanced or recurrent lung cancer, it aims to control disease and symptoms. Your team should mark which goal applies to you; benefit is not guaranteed.

Your treatment plan

A cycle lasts 21 days (3 weeks). Vinorelbine is given on Days 1, 8, 15. Cisplatin is given on Day 1. Four cycles are planned, with review before doses. Delays or omitted doses may be needed for safety.

1. **Before treatment**

Confirm your treatment goal, blood tests and contact numbers.

2. **Four planned cycles**

A cycle lasts 21 days (3 weeks). Vinorelbine is given on Days 1, 8, 15. Cisplatin is given on Day 1. Four cycles are planned, with review before doses. Delays or omitted doses may be needed for safety.

3. **After treatment**

Your team reviews recovery and agrees scans and any further treatment.

Before and between visits

- Bring your medicines list and SACT card.
- Have blood tests as arranged before treatment.
- Tell the team about bowel, hearing or nerve changes.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately for temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions.

Cannot reach the team? Seek urgent medical assessment. Do not wait for the next appointment.

Your contact plan

Ask your team to complete these before you leave.

Hospital / team _____

Daytime number and hours _____

Evening / weekend number _____

If unavailable, where to go _____

Keep this with your consent information. Ask your team about anything unclear.

What to watch for

Use the picture and symptom descriptions together. You do not need to know which medicine caused a symptom before calling.

1. Hearing and nerves

Cisplatin / vinorelbine

Report ringing ears, hearing change, numbness or weakness; effects can persist.

2. Mouth

Both medicines

Report mouth sores that make eating or drinking difficult.

3. Kidneys / fluids

Cisplatin

Fluids protect kidneys; call for very little urine or inability to drink.

4. Bowels / sickness


Both medicines

Constipation and nausea need early treatment; severe pain or vomiting needs urgent advice.

5. Blood cells / infusion site

Both medicines

Fever or sudden illness needs immediate advice. Tell the nurse about infusion pain.



Selected effects, not every possible effect or a prediction. Match each number to its symptom box. Blood, skin and nerve markers are schematic.

Call your oncology team immediately for fever or sudden illness, shaking chills, breathing difficulty, chest pain, uncontrolled vomiting, very little urine, severe abdominal pain or constipation with vomiting. Tell the nurse immediately about burning or pain at the infusion site. Do not wait until the next visit.

My SACT card

Vinorelbine and cisplatin · 21-day cycle

21-day cycle. Vinorelbine Days 1, 8, 15; cisplatin Days 1.

Fever or sudden illness: call oncology immediately. Report reduced urine, severe constipation or infusion-site pain.

Name: _____
Hospital: _____
Oncology phone: _____
Out of hours: _____

Show this card when seeking care. It does not replace a medicine-specific alert card.

More information, when you want it

Scan for medicines, tests, daily life and your treatment record.

https://sactcheck.com/patient/00339/