BASELINE → TREATMENT → FOLLOW-UP

Three treatment visits and an interval

This 28-day schedule has no Day 22 dose. No fixed scan interval or guaranteed recovery day.

Check	Before starting	Day1	Day8	Day15	Days22–28	Nextcycle
Paclitaxel	Confirm aim and off-label consent	Infusion	Infusion if appropriate	Infusion if appropriate	No Day 22 dose	Repeat if helping/ tolerated
Tests and symptoms	FBC, kidney, liver; nerves	Review before dose	Review before dose	Review before dose	Call if unwell	Review before dose
Premedication and review	Confirm first-dose plan	Prescribed premedication	Later-dose plan	Later-dose plan	Continue symptom plan	Confirm appointments

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00621 v3 · official regimen

THE WHOLE TREATMENT JOURNEY

Paclitaxel Days 1,8 and 15 · relapsed or refractorySCLC: treatment and checks

Lung · relapsed/refractorysmall-cell lung cancer

Paclitaxel is given on Days 1,8 and 15 of each 28-day cycle. Day 22 is a planned week off, not a fourth infusion. Treatment continues while helping and tolerated; there is no fixed cycle limit.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Day1	Day8	Day15	Days22–28	Nextcycle
Paclitaxel NCCP 00621	Confirm aim and off-label consent	Infusion	Infusion if appropriate	Infusion if appropriate	No Day 22 dose	Repeat if helping/ tolerated
Tests and symptoms NCCP 00621	FBC, kidney, liver; nerves	Review before dose	Review before dose	Review before dose	Call if unwell	Review before dose
Premedication and review NCCP 00621	Confirm first-dose plan	Prescribed premedication	Later-dose plan	Later-dose plan	Continue symptom plan	Confirm appointments

What each check is for

1 Before each dose

Days 1,8 and 15

Blood results and symptoms can change

Ask / act: Review before every dose and confirm premedication.

Sources: NCCP 00621

2 Fourthweek

Days 22–28

No scheduled infusion

Ask / act: Call if unwell; agree the next cycle and response-review plan.

Sources: NCCP 00621

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.28