SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Carboplatin + Oral Etoposide

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Extensive-disease small-cell lung cancer · carboplatin + oral etoposide · NCCP 00319 · version 5 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
This chemotherapy combination aims to control extensive-disease small-cell lung cancer, ease symptoms and help people live longer. Expected benefit depends on the cancer, general health and treatment choices. Ask what the team hopes it will achieve for you.

Schedule
Carboplatin is planned into a vein on Day 1, before oral etoposide. Etoposide capsules are prescribed twice daily on Days 1, 2 and 3 only, on an empty stomach. The cycle repeats every 21 days while treatment is helping and side effects remain acceptable. Your team decides when to delay, change or stop; the source does not set a fixed number of cycles.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Carboplatin
PLATINUM CHEMOTHERAPY

- ☐ Low blood counts, infection and bleeding
- ☐ Nausea, vomiting and fatigue
- ☐ Mouth soreness and diarrhoea
- ☐ Kidney problems
- ☐ Allergic or infusion reactions
- ☐ Hearing changes or tinnitus, which can persist
- ☐ Numbness or tingling, which can persist

Oral etoposide
TOPOISOMERASE-II INHIBITOR CHEMOTHERAPY

- ☐ Low blood counts, infection and bleeding
- ☐ Nausea, vomiting, diarrhoea and mouth soreness
- ☐ Hair loss and fatigue
- ☐ Allergic reactions
- ☐ Fertility effects
- ☐ Rare serious skin reactions
- ☐ Secondary leukaemia can occur

Discuss the individual treatment aim and likely benefit, uncertainty, alternatives including not proceeding, common and serious risks, lasting hearing/nerve/kidney effects, fertility and pregnancy advice, safe capsule use, practical support and the possibility of dose delays or changes. Oral treatment is chemotherapy and needs the same careful safety review as an infusion.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00319 v5.

SACTCheck v0.84.23
Developed by Dr Paul O'Brien