NCCP 00908 · clinician draft

Serplulimab, carboplatin and etoposide

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Maintenance: serplulimab alone

Active medicines: serplulimab

Emetogenic risk: minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Serplulimab alone has minimal emetogenic potential: no routine continuation of the induction chemotherapy antiemetic course.
- Provide symptom-directed rescue if appropriate; investigate new nausea/vomiting for endocrine, immune and other causes.
- Continue cycle bloods, thyroid tests every3–6weeks and immune symptom vigilance; keep the alert card.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00908 v1b · p7–8 · NCCP NCIS Medical Oncology antiemetics V8

Sources checked 2026-10-26 · SACTCheck v0.84.23 · Regimen-level drafts; prescriber and oncology-pharmacy verification required