

Adjuvant medication prescription for chemotherapy Moderate Risk for nausea and vomiting

Name			Allergies	
DOB				
MRN			Date of prescription	

- Dexamethasone 8mg OD PO on days 2&3 of chemotherapy**
Directions: Take 8mg daily for two days starting the day after chemotherapy.
- Omeprazole 20mg OD PO on days 1-3 of chemotherapy**
Directions: Take one capsule daily for three days starting the day of chemotherapy.
- Nystatin 100,000units/mL 1ml QDS PO continuous during chemotherapy**
Directions: Rinse 1mL around the mouth four times a day after meals during chemotherapy regimen.
- Chlorhexidine mouthwash 10ml QDS PO continuous during chemotherapy**
Directions: Rinse 10mls around the mouth for at least 1 minute four times a day during chemotherapy regimen. The mouthwash should be expelled from the mouth after rinsing.
- Cyclizine 50mg TDS PO PRN**
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- Metocloperamide 10mg TDS PO PRN**
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- Loperamide 4mg STAT then 2mg QDS PO PRN, (max 16mg/day) for chemotherapy-induced diarrhoea**
Directions: Take two tablets after first episode of diarrhoea, then one after every diarrheal episode

Repeat by: ____ cycles
(cycle length = ____ weeks)

Signed:	
Name (print):	
IMC number:	

Adjuvant medication prescription for chemotherapy Moderate Risk for nausea and vomiting

Name			Allergies	
DOB				
MRN			Date of prescription	

- Dexamethasone 8mg OD PO on days 2&3 of chemotherapy**
Directions: Take 8mg daily for two days starting the day after chemotherapy.
- Omeprazole 20mg OD PO on days 1-3 of chemotherapy**
Directions: Take one capsule daily for three days starting the day of chemotherapy.
- Nystatin 100,000units/mL 1ml QDS PO continuous during chemotherapy**
Directions: Rinse 1mL around the mouth four times a day after meals during chemotherapy regimen.
- Chlorhexidine mouthwash 10ml QDS PO continuous during chemotherapy**
Directions: Rinse 10mls around the mouth for at least 1 minute four times a day during chemotherapy regimen. The mouthwash should be expelled from the mouth after rinsing.
- Cyclizine 50mg TDS PO PRN**
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- Metocloperamide 10mg TDS PO PRN**
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- Loperamide 4mg STAT then 2mg QDS PO PRN, (max 16mg/day) for chemotherapy-induced diarrhoea**
Directions: Take two tablets after first episode of diarrhoea, then one after every diarrheal episode

Repeat by: ____ cycles
(cycle length = ____ weeks)

Signed:	
Name (print):	
IMC number:	