

Trifluridine/tipiracil (Lonsurf)

Gastrointestinal · NCCP 00382 · version 4

Your medicines

Your treatment schedule

A picture of side effects

Numbers connect each symptom box to the relevant anatomical area. Tell your team about new or worsening symptoms.

Trifluridine/tipiracil · oral chemotherapy

1 Blood / infection

Trifluridine/tipiracil

Low white cells can increase infection risk. Fever, shivers or sudden illness needs immediate contact.

2 Mouth

Trifluridine/tipiracil

Report mouth soreness or ulcers that limit eating or drinking.

3 Stomach / bowel

Trifluridine/tipiracil

Nausea, vomiting or diarrhoea can cause dehydration if persistent.

4 Energy / fatigue

Trifluridine/tipiracil

Report sudden or marked tiredness; infection, anaemia or dehydration can also contribute.

5 Skin


Trifluridine/tipiracil

Report persistent or severe rash or other skin changes.

6 Tablet handling

Your prescribed tablets

Follow the pharmacy handling and storage instructions. Keep tablets in the original packaging.



Selected effects only: the same symptom can have more than one cause, and not every possible side effect is shown.

Trifluridine/tipiracil (Lonsurf)

Gastrointestinal · NCCP 00382 · version 4

TREATMENT JOURNEY

Days 1-5 Take the prescribed dose twice daily.	Days 6-7 No Lonsurf.	Days 8-12 Take the prescribed dose twice daily.	Days 13-28 No Lonsurf; recovery and monitoring period.	Next cycle A new 28-day cycle starts only if review and blood tests allow.
--	--------------------------------	---	--	--

IMPORTANT MEDICINE-SPECIFIC EFFECTS

Trifluridine/tipiracil (Lonsurf)

- Low neutrophils, anaemia and platelets can require interruption or dose reduction.
- Fever with neutropenia needs urgent assessment.
- Nausea, vomiting and diarrhoea can cause dehydration.

OTHER / SHARED TREATMENT EFFECTS

The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed.

Monitoring / medicines

- Take doses after the morning and evening meals exactly as prescribed.
- Do not double a dose for a missed or vomited dose; follow your oncology/pharmacy instructions.
- Blood counts are particularly important because neutropenia can be significant.

Bring / track

- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT

Call oncology urgently for fever, shivers, sudden illness, uncontrolled vomiting, persistent diarrhoea, bleeding, inability to drink or marked weakness. Severe breathing difficulty, collapse or severe chest pain: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00382/>

Official NCCP 00382 v4

SACTCheck v0.81.2 · patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

