

Nivolumab + ipilimumab + pemetrexed + cisplatin

Lung · NCCP 00714 · version 4

Your medicines

Your treatment schedule

A picture of side effects

Numbers connect each symptom box to the relevant anatomical area. Tell your team about new or worsening symptoms.

Nivolumab / ipilimumab · immune effects

Pemetrexed / cisplatin · chemotherapy

6 Hearing / nerves

Cisplatin

Report tinnitus, hearing change, numbness or tingling before it worsens.

3 Heart / muscles

Nivolumab / ipilimumab

Chest pain, palpitations, fainting or marked muscle weakness needs urgent assessment.

5 Kidneys / hydration

Pemetrexed / cisplatin

Kidney function is monitored closely and cisplatin requires hydration/electrolyte support.

7 Blood / infection

Pemetrexed / cisplatin

Fever, shivers, bruising or sudden illness can reflect low blood counts.

2 Lungs

Nivolumab / ipilimumab

New cough or breathlessness needs prompt assessment for pneumonitis and other causes.

4 Liver / bowel

Nivolumab / ipilimumab

Jaundice, dark urine, persistent diarrhoea or abdominal pain can indicate immune inflammation.

6 Hearing / nerves


Cisplatin

Report tinnitus, hearing change, numbness or tingling before it worsens.

8 Nausea / mouth

Cisplatin / pemetrexed

Use the prescribed anti-sickness medicines; persistent vomiting or severe mouth soreness needs advice.



Selected effects only: the same symptom can have more than one cause, and not every possible side effect is shown.

Nivolumab + ipilimumab + pemetrexed + cisplatin

Lung · NCCP 00714 · version 4

TREATMENT JOURNEY

Cycle 1 · day 1	Cycle 1 · day 22	After cycle 1	Duration
Nivolumab + ipilimumab + pemetrexed + cisplatin.	Nivolumab + pemetrexed + cisplatin.	Continue nivolumab on days 1 and 22 and ipilimumab on day 1 of each 42-day cycle.	Immunotherapy continues until progression/unacceptable toxicity or up to 24 months in patients without progression, according to the NCCP

IMPORTANT MEDICINE-SPECIFIC EFFECTS

Nivolumab + ipilimumab - immune effects

- Immune inflammation: lungs, bowel, liver, hormone glands, kidneys, skin, heart, nerves or muscles.
- New immune symptoms can begin during treatment or after treatment has stopped.

Pemetrexed

- Low blood counts can increase infection, anaemia or bruising risk.
- Mouth/gut inflammation and skin rash can occur.
- Kidney function can deteriorate and is monitored.

Cisplatin

- Kidney injury and low magnesium/potassium can occur.
- Hearing loss or ringing in the ears may be permanent.
- Numbness/tingling and marked nausea/vomiting can occur.

OTHER / SHARED TREATMENT EFFECTS

The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed.

Monitoring / medicines

- Folic acid and vitamin B12 support are required with pemetrexed.
- Cisplatin requires hydration/electrolyte management and high-emetogenic-risk antiemetic support.
- Blood count, kidney/liver profile and endocrine monitoring support treatment safety.

Bring / track

- Immune adverse events can begin during treatment or after it stops; report new symptoms early.

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT

Contact oncology urgently for fever, new breathlessness, persistent diarrhoea, jaundice, severe rash, chest pain, marked weakness, inability to maintain fluids, reduced urine or new neurological symptoms. For severe breathing difficulty, collapse, seizure or severe chest pain call 112/999.

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00714/>

Official NCCP 00714 v4

SACTCheck v0.81.2 · patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

