

Atezolizumab + bevacizumab

Advanced or unresectable hepatocellular carcinoma / NCCP 00831 v2a

Your medicines

Your treatment schedule

A picture of side effects

Numbers connect each symptom box to the relevant anatomical area. Tell your team about new or worsening symptoms.

Atezolizumab · immune effects

Bevacizumab · vessels / healing

1 Hormones / brain

Atezolizumab

Tiredness, thirst, dizziness or a new headache (hormone or pituitary inflammation).

3 Heart

Atezolizumab

Chest pain, palpitations or shortness of breath (myocarditis / pericarditis).

5 Kidneys

Atezolizumab

Passing less urine, swelling or blood test changes (nephritis). Bevacizumab can also cause protein in the urine, usually found on a urine test.

7 Skin / nerves / muscles

Atezolizumab

Rash or blisters; weakness, numbness or muscle pain (skin inflammation / nerve effects / myositis).

9 Blood vessels / healing

Bevacizumab

Bleeding, clots or slow wound healing (bevacizumab-related effects).

2 Lungs

Atezolizumab

New cough or shortness of breath (pneumonitis).

4 Liver

Atezolizumab

Yellow eyes, dark urine or changes in blood tests (hepatitis).

6 Bowel


Atezolizumab

New diarrhoea, blood or tummy pain (colitis; severe pain may have another urgent cause). Bevacizumab can rarely cause a gut perforation; severe belly pain

8 Joints

Atezolizumab

Painful, swollen joints (inflammatory arthritis).



Selected effects only: the same symptom can have more than one cause, and not every possible side effect is shown.

Atezolizumab + bevacizumab

Advanced or unresectable hepatocellular carcinoma / NCCP 00831 v2a

TREATMENT JOURNEY

Before C1	C1 · Day 1	Days 2–21	C2 · Day 22	Later / after
Varices screening; blood count, liver/kidney, glucose, thyroid, BP and urine protein assessment.	Atezolizumab IV or SC, then bevacizumab IV. Tell the nurse immediately about an infusion or injection reaction.	Notice new symptoms. Follow any home BP instructions and keep your SOS number accessible.	Repeat review, bloods, BP and urine; thyroid testing is scheduled every six weeks. Treatment only if appropriate.	Each cycle follows the same 21-day pattern. Immune effects may still appear after treatment stops.

IMPORTANT MEDICINE-SPECIFIC EFFECTS

Atezolizumab - immune effects

- Immune inflammation: lungs, bowel, liver, hormone glands, kidneys, skin, heart, nerves or muscles.
- New immune symptoms can begin during treatment or after treatment has stopped.

Bevacizumab

- Blood pressure and urine protein can rise.
- Bleeding, blood clots and impaired wound healing can occur.
- Rare bowel perforation/fistula or severe neurological symptoms (PRES) need urgent assessment.

OTHER / SHARED TREATMENT EFFECTS

The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed.

Monitoring / medicines

- Before starting Bleeding and liver context Oesophageal varices screening, BP, urine protein, blood count, liver/kidney profile, glucose and thyroid tests. Discuss past
- Each cycle Blood, urine, BP NCCP specifies blood count, liver/kidney profile, glucose and urine-protein dipstick before each cycle, plus BP before and after treatment. Results
- Every 6 weeks Thyroid tests Thyroid function tests are scheduled every six weeks. Your team may order additional endocrine tests when symptoms suggest adrenal, pituitary or

Bring / track

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT

04 / action language Know when to call Emergency: call 112/999 Collapse, severe bleeding or vomiting blood, severe chest pain, severe breathing difficulty, seizure, sudden new vision loss or a major neurological change. Tell responders you receive atezolizumab and bevacizumab. Contact oncology urgently Black stools, any concerning bleeding, severe or persistent diarrhoea, new cough or breathlessness, yellow eyes, reduced urine, new severe headache, new vision changes, marked weakness, confusion, a temperature $\geq 37.5^{\circ}\text{C}$ or $\leq 36^{\circ}\text{C}$, or feeling suddenly unwell. Follow your centre's SOS instructions even if a symptom begins after the last cycle. Medicines and fever Measure

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00831/>

Official NCCP 00831 v2a

SACTCheck v0.81.2 · patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

