SACTCHECK TREATMENT ASSESSMENT

Bevacizumab + mFOLFOX6

NCCP 00446 | 7b
SACTCheck v0.83.3

TUMOUR GROUP	INDICATION	COURSE
Gastrointestinal	**Treatment of adult patients with metastatic carcinoma of the colon or rectum.**	**Course metadata not confirmed**

TREATMENT CONTEXT	ASSESSMENT ID	ASSESSMENT TIME
Not entered	**JSON-DEMO-001**	**16/10/2026, 22:21**

ENCODED CRITERIA RESULT

1 encoded criterion requires clinical review

Review the highlighted protocol criteria and the current NCCP source before the final treatment decision.

ENTERED VALUES AND ENCODED CRITERIA

Modified FOLFOX-6 assessment

DOMAIN	ENTERED VALUE	ENCODED CRITERION	RESULT
Proteinuria (g/24 h)	4.1	> 4	**Criterion not met - urgent review**

COMPONENT-SPECIFIC PROTOCOL ACTIONS

Protocol-level guidance; verify prescribed dose/current level

Bevacizumab: Discontinue - Not to be administered
Discontinue bevacizumab when 24-hour urine protein exceeds 4 g/24 h. page 5 | Table 4

UNASSESSED DOMAINS

Known hypersensitivity to regimen component or excipient, Pregnant, Breastfeeding, Assessment type, Baseline peripheral neuropathy with functional impairment, DPD status, ANC, Platelets + 26 more

Blank domains were not assumed normal.

CLINICIAN DECISION

Not documented
Not documented

CLINICAL DECISION SUPPORT - NOT TREATMENT CLEARANCE

SACTCheck applies encoded NCCP protocol criteria to clinician-entered information. This output does not constitute treatment clearance and does not replace complete clinical assessment, the current official NCCP protocol, prescribing information, local policy or professional judgement. The responsible oncology clinician retains responsibility for the final treatment decision.