
Panitumumab

Day 1: panitumumab infusion. There is no home chemotherapy pump in this regimen. A cycle lasts 14 days. Treatment continues while it is helping and side effects remain acceptable.

Draft for clinical, pharmacy and patient/PPI review. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.



1 Skin / rash

Panitumumab

Report rash, cracks, painful or infected skin early.

2 Eyes

Panitumumab

New eye pain, light sensitivity or blurred vision: prompt advice.

3 Breathing

Panitumumab

New or worsening breathlessness or cough needs assessment.

4 Bowel

Panitumumab

Severe diarrhoea, dizziness or inability to drink: urgent contact.

5 Nails

Panitumumab

Painful or infected nail folds need early treatment.

6 Blood minerals

Panitumumab

Tests check magnesium, calcium and potassium; report cramps or weakness.

7 Hydration

Panitumumab

Reduced urine or severe dehydration needs urgent advice.

8 Infusion reaction

Panitumumab

Tell the nurse about swelling, itching, flushing or breathing changes.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Panitumumab

- Rash, dry skin, cracks and sore nails
- Eye inflammation or corneal injury
- Low magnesium, calcium or potassium
- Diarrhoea, dehydration and kidney problems
- Infusion reaction; rare serious lung or skin effects

Make the plan practical

Start skin care early

Ask for your skin-care plan before the first infusion. Use the recommended moisturiser and sun protection; your team may prescribe a cream or an antibiotic. Do not use ordinary acne treatments without advice.

Minerals and blood tests

Panitumumab can lower magnesium and other minerals even when you feel well. Blood tests continue during treatment and for about eight weeks after it finishes. Take replacement treatment only as prescribed.

Feeling sick or loose stools

Panitumumab has low nausea risk, but symptoms can still occur. Follow the rescue medicine and hydration plan from your team. Severe or persistent diarrhoea, reduced urine or inability to drink needs urgent advice.

Travel and daily life

Allow time for blood tests, review, preparation, infusion and observation. Ask about transport if medicines or symptoms affect driving. Pace activity and seek help with food, work or caring responsibilities.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

CUH · MON–FRI 8AM–5PM

087 181 9645

Acute Oncology Service

CUH · WEEKENDS / BANK HOLIDAYS 9AM–5PM

021 4922000

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00225/

Sources and review status are available online. Follow your own prescription and hospital instructions.