SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Panitumumab + FOLFIRI

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

RAS wild-type metastatic colorectal cancer · NCCP 00448 · version 4b · 14-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Panitumumab + FOLFIRI is used for bowel cancer that has spread and has a confirmed wild-type RAS result. This NCCP regimen covers first-line treatment and selected second-line treatment after a fluoropyrimidine regimen without irinotecan. Your team explains the aim, alternatives and how benefit will be reviewed.

Schedule
Day 1: panitumumab, irinotecan, folinic acid and fluorouracil, followed by a fluorouracil pump for about 46 hours; disconnection is arranged around Day 3. A cycle lasts 14 days. Treatment continues while it is helping and side effects remain acceptable.

1. GENERIC SACT RISKS / DISCUSSION POINTS

- ☐ Purpose, expected benefit and treatment plan explained
- ☐ Treatment may be delayed, modified or stopped because of toxicity
- ☐ Unexpected, rare or serious adverse effects can occur
- ☐ Supportive medicines and urgent contact advice discussed
- ☐ Pregnancy avoidance / contraception / fertility discussed where relevant
- ☐ Bone-marrow suppression / infection risk
- ☐ Anaemia / fatigue and low platelets / bleeding
- ☐ Nausea / vomiting and bowel disturbance
- ☐ Mouth soreness / mucositis and skin changes
- ☐ Allergic / infusion reaction and other organ-specific toxicity

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Panitumumab
EGFR-TARGETED ANTIBODY

- ☐ Rash, dry skin, cracks and sore nails
- ☐ Eye inflammation or corneal injury
- ☐ Low magnesium, calcium or potassium
- ☐ Diarrhoea, dehydration and kidney problems
- ☐ Infusion reaction; rare serious lung or skin effects

Irinotecan
TOPOISOMERASE I CHEMOTHERAPY

- ☐ Early cramps, sweating and diarrhoea
- ☐ Delayed diarrhoea and dehydration
- ☐ Low blood counts / serious infection
- ☐ Nausea, mouth soreness and fatigue
- ☐ Hair thinning or loss
- ☐ Rare serious lung or allergic reactions

Folinic acid
CHEMOTHERAPY MODULATOR

- ☐ Enhances fluorouracil activity and toxicity
- ☐ Rare allergic reaction
- ☐ Diarrhoea / mouth soreness in combination

Fluorouracil
CHEMOTHERAPY

- ☐ Diarrhoea, dehydration and mouth ulcers
- ☐ Low blood counts / infection
- ☐ Hand-foot skin soreness
- ☐ Cardiotoxicity including chest pain / ischaemia
- ☐ Severe early toxicity including DPD-related toxicity
- ☐ Pump / central-line complications

Discuss RAS selection, skin/nail and eye effects, electrolyte monitoring for eight weeks after completion, infusion reactions and rare serious lung/skin toxicity. Confirm DPD testing, antiemetics, atropine suitability and the separate delayed-diarrhoea plan. Explain port/PICC care and pump connection/disconnection.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00448 v4b.

SACTCheck v0.83.5
Developed by Dr Paul O'Brien