SACTCHECK TREATMENT ASSESSMENT

Panitumumab 6 mg/kg Therapy

NCCP 00225 | 6
SACTCheck v0.83.5

TUMOUR GROUP	INDICATION	COURSE
Gastrointestinal	**RAS wild-type metastatic colorectal cancer after fluoropyrimidine, oxaliplatin and irinotecan.**	**Advanced disease | q14d**

TREATMENT CONTEXT	ASSESSMENT ID	ASSESSMENT TIME
Not entered	**JSON-DEMO-001**	**17/10/2026, 00:22**

ENCODED CRITERIA RESULT

1 encoded criterion requires clinical review

Review the highlighted protocol criteria and the current NCCP source before the final treatment decision.

ENTERED VALUES AND ENCODED CRITERIA

Standard assessment

DOMAIN	ENTERED VALUE	ENCODED CRITERION	RESULT
Current panitumumab skin toxicity grade	Grade 1 - Mild	< 3	**Dose-modification pathway**
Doses withheld for this skin episode	1	in 1, 2	**Dose-modification pathway**
Not recovered after the planned 1-2 dose withholding period	No	= No	**Dose-modification pathway**
Number of episodes of grade >= skin toxicity	2	= 2	**Dose-modification pathway**

COMPONENT-SPECIFIC PROTOCOL ACTIONS

Protocol-level guidance; verify prescribed dose/current level

Panitumumab: Use 80% of protocol dose - 4.8 mg/kg
After recovery below grade 3 following 1-2 withheld doses: panitumumab 80% of original dose. Confirm suitability and previous individual modifications. page 2 | Table 2

UNASSESSED DOMAINS

Validated tumour RAS status, ECOG performance status, Known hypersensitivity to a regimen component, Confirmed interstitial pneumonitis / pulmonary fibrosis, New or worsening unexplained respiratory symptoms, Severe skin infection, blistering or skin detachment, Infusion-related reaction grade, New eye pain, redness, light sensitivity or blurred vision + 7 more

Blank domains were not assumed normal.

CLINICIAN DECISION

Not documented
Not documented

CLINICAL DECISION SUPPORT - NOT TREATMENT CLEARANCE

SACTCheck applies encoded NCCP protocol criteria to clinician-entered information. This output does not constitute treatment clearance and does not replace complete clinical assessment, the current official NCCP protocol, prescribing information, local policy or professional judgement. The responsible oncology clinician retains responsibility for the final treatment decision.