SACTCHECK TREATMENT ASSESSMENT

Panitumumab + mFOLFOX6

NCCP 00447 | 6b
SACTCheck v0.83.5

TUMOUR GROUP	INDICATION	COURSE
Gastrointestinal	**First-line treatment of adult patients with wild-type RAS metastatic colorectal cancer.**	**Course metadata not confirmed**

TREATMENT CONTEXT	ASSESSMENT ID	ASSESSMENT TIME
Not entered	**JSON-DEMO-001**	**17/10/2026, 00:23**

ENCODED CRITERIA RESULT

1 encoded criterion requires clinical review

Review the highlighted protocol criteria and the current NCCP source before the final treatment decision.

ENTERED VALUES AND ENCODED CRITERIA

Standard assessment

DOMAIN	ENTERED VALUE	ENCODED CRITERION	RESULT
Peripheral neuropathy grade	Grade 2	= 2 CTCAE grade	**Dose-modification pathway**
Current oxaliplatin dose level	Level -1	Linked context required	**Additional clinical review/context**

COMPONENT-SPECIFIC PROTOCOL ACTIONS

Protocol-level guidance; verify prescribed dose/current level

Oxaliplatin: Dose level -2 - 50 mg/m^2
Grade 2 neuropathy at the start of a cycle requires a one-level oxaliplatin reduction. page 5 | Table 5

UNASSESSED DOMAINS

Validated tumour RAS status, ECOG performance status, Known hypersensitivity to regimen component or excipient, Confirmed interstitial pneumonitis / pulmonary fibrosis, New or worsening unexplained respiratory symptoms, Current panitumumab skin toxicity grade, Number of episodes of grade >=3 skin toxicity, Not recovered after the planned 1-2 dose withholding period + 30 more

Blank domains were not assumed normal.

CLINICIAN DECISION

Not documented
Not documented

CLINICAL DECISION SUPPORT - NOT TREATMENT CLEARANCE

SACTCheck applies encoded NCCP protocol criteria to clinician-entered information. This output does not constitute treatment clearance and does not replace complete clinical assessment, the current official NCCP protocol, prescribing information, local policy or professional judgement. The responsible oncology clinician retains responsibility for the final treatment decision.