SACTCHECK · PATIENT TREATMENT GUIDE

Durvalumab monotherapy

NCCP 00655 · v3a

Your medicines

durvalumab

Your treatment schedule


Phase 1
28-day cycle
Day 1 durvalumab (IV infusion)

Chemo man guide

Where side effects may appear

Use this simple body map to explain that treatment effects can involve different body systems. The exact important effects for this regimen are shown beside it.

Head / hormones **Lungs** **Skin** **Gut** **Liver** **Kidneys / urine** **Nerves / muscles**



Immunotherapy-related effects

New cough or breathlessness (pneumonitis)
New or worsening cough, breathlessness, chest discomfort or reduced exercise tolerance.
Durvalumab

Diarrhoea or bowel inflammation (colitis)
New diarrhoea, abdominal pain, blood or mucus in the stool.
Durvalumab

Liver inflammation (hepatitis)
Yellowing of the skin/eyes, dark urine, nausea or abnormal liver blood tests.
Durvalumab

Hormone changes (endocrine problems)
Unusual fatigue, dizziness, headache, thirst, frequent urination, temperature intolerance or weakness.
Durvalumab

Kidney inflammation (nephritis)
Reduced urine, swelling, nausea or changes in kidney blood tests.
Durvalumab

New rash or blisters (skin toxicity)
New or worsening rash; urgent review for blistering, painful skin or mouth involvement.
Durvalumab

Contact your oncology team urgently for severe or rapidly worsening symptoms, fever or sudden illness, new breathlessness or chest pain, persistent diarrhoea or vomiting, neurological change, significant bleeding, or another symptom your team has told you is urgent.

More detailed regimen information

Scan to reopen this regimen guide. The QR contains a public regimen link and no patient data.


https://sactcheck.com/?patientSupport=nccp-00655-v3a

Patient information support only. This does not replace your oncology team's advice, formal consent, current NCCP guidance or local emergency instructions.

THE WHOLE TREATMENT COURSE

Chemoradiotherapy first, then durvalumab

For this lung-cancer pathway, chemotherapy and radiotherapy are given together. Durvalumab is a separate later phase, after treatment review.



The horizontal axis shows order, not equal time intervals. NCCP specifies starting durvalumab within 42 days after chemoradiotherapy and continuing for up to 12 months (13 cycles), if suitable. The team confirms recovery, no progression and all eligibility requirements.

> **Chemotherapy and radiotherapy overlap in the earlier phase. Durvalumab follows; it is not shown as concurrent with them.**

Educational reference, not a personal appointment calendar. NCCP 00655 v3a, pages 1–2