Enzalutamide: castration-resistant prostate cancer

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Continuous oral treatment

Active medicines: Enzalutamide

Emetogenic risk: Minimal to low (oral)

Medicine	Dose / route	When
Antiemetic plan for current oral dose	Individual prescription	As agreed with oncology/pharmacy Review QT prolongation and all interactions; no automatic intravenous (IV) chemotherapy script.

Additional phase-specific support

- Take at the same time daily with or without food. Swallow the prescribed capsules or tablets whole with water; do not open, chew or crush them. If missed, take as soon as remembered that day. If the entire day is missed, resume the usual dose the following day; do not double it. Ask pharmacy if swallowing is difficult.
- Blood count, kidney and liver tests every 4 weeks as clinically indicated. Blood pressure, electrocardiogram (ECG) and creatinine as clinically indicated. If taking warfarin, international normalised ratio (blood-clotting test) (INR) weekly until a stable dose is established.
- Report falls, new memory or concentration problems and troublesome fatigue. Seizure, severe headache with confusion or vision changes needs urgent help. Discuss driving safety and every new medicine with the team; interactions can change anticoagulant effects.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00233 v7 · pTreatment / supportive care · NCCP regimen and NCIS antiemetic guidance

Sources checked 2026-11-14 · SACTCheck v0.84.51 · Regimen-level drafts; prescriber and oncology-pharmacy verification required