Apalutamide

To control prostate cancer and delay progression; your team explains the expected benefit for your situation.

High-risk nonmetastatic castration-resistant or metastatic hormone-sensitive prostate cancer · NCCP 00574 v5 · Document revised 14 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

High-risk nonmetastatic castration-resistant or metastatic hormone-sensitive prostate cancer

My overall treatment

Once daily plus ongoing ADT

28 day reference

Reference window · ongoing treatment

Apalutamide every day — Day 1–28

Treatment

Apalutamide 240 mg once daily, continuously while helping and tolerated, with ongoing ADT unless surgically castrated. Lower prescribed doses are 180 mg or 120 mg daily when indicated.

Review

Continue ADT. The team reviews symptoms, blood tests, PSA and scans and confirms whether treatment continues.

Your 28-day review window

Day 1–28

Every day, including weekends

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline preparation** NCCP	Full blood count, kidney and liver tests and thyroid function; electrocardiogram (ECG) if at risk of heart-rhythm changes; international normalised ratio (blood-clotting test) (INR) if taking warfarin. Review cardiovascular and seizure history, falls, bone health and all medicines.
During	**Safety checks** NCCP	Repeat blood count, kidney and liver tests on the schedule agreed with your team. Thyroid function and ECG as clinically indicated. If warfarin is unavoidable, INR weekly until the dose is stable. The NCCP regimen does not set one fixed interval for every routine blood test.
Response	**Cancer-specific checks** Individual specialist plan	PSA and testosterone assess cancer activity and hormone suppression; scans follow your individual plan.

Between treatments

- Symptoms can occur on any day; keep your urgent contact plan.
- Report rash promptly. Blisters, peeling skin, fever with rash, or mouth/eye sores need immediate medical advice. New or worsening cough or breathlessness requires urgent assessment for lung inflammation. Seizures require emergency help. Discuss driving and medicine interactions.

Bring your medicines list, calendar and verified contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Energy, balance and thinking

Apalutamide

Report troublesome fatigue, falls or confusion. Seizure, collapse or sudden neurological symptoms need emergency help.

3 Heart and blood pressure

Apalutamide

Keep blood-pressure checks. Chest pain, fainting or severe breathlessness needs urgent help; report ankle swelling or palpitations.

5 Breathing and lung inflammation


Apalutamide

New or worsening cough or breathlessness needs urgent assessment. Treatment-related lung inflammation can be severe.

7 Sexual health and fertility

ADT and Apalutamide

Discuss erections, libido, fertility and contraception. Avoid exposing a pregnant partner to these medicines; follow the product-specific precautions.



2 Thyroid function

Apalutamide

Keep thyroid blood tests. An underactive thyroid can cause fatigue, weight changes or feeling cold; report new symptoms.

4 Liver and blood tests

Apalutamide

Keep liver blood tests. Report yellow skin or eyes, dark urine, reduced appetite or a major change in wellbeing.

6 Skin changes

Apalutamide

Report a new rash. Severe rash with blisters, peeling, fever or mouth/eye sores needs urgent assessment.

8 Bones and muscle strength

ADT and Apalutamide

Discuss bone-density assessment, falls prevention, suitable resistance and aerobic activity, calcium intake and vitamin D needs.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology urgently for sudden illness, persistent vomiting, inability to drink, jaundice, new breathlessness or severe rash. Call 112/999 for severe chest pain, severe breathing difficulty, collapse, seizure, or new leg weakness with loss of bladder or bowel control.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

High-risk nonmetastatic castration-resistant or metastatic hormone-sensitive prostate cancer · NCCP 00574 v5 · Document revised 14 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Apalutamide

Once daily plus ongoing ADT

Phase: _____ Continuous treatment: team reviews dose

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My review record

Recording spaces only; not a fixed course.

Review	Date / initials

My treatment information

Latest review date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team records each review and the current daily dose. Recording boxes do not imply a fixed course or a rest period. Record any held dose or changed plan. An entry does not authorise a restart; keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.