

Pembrolizumab + pemetrexed + carboplatin

Lung · NCCP 00568 · version 5

Your medicines


Pembrolizumab · Pemetrexed · Carboplatin

Your treatment schedule

Every 21 days; initial combination followed by maintenance when appropriate.

A picture of side effects

Tell your team about new or worsening symptoms. The same symptom can have more than one cause. Numbers connect each symptom box to its location.



This map shows selected effects to discuss; it does not show their likelihood or every possible side effect. Blood-count and whole-body effects are marked schematically.

Selected symptoms to help discussion. Tell your team about any new or worsening symptom. See page 2 for the schedule, support and QR.

Pembrolizumab + pemetrexed + carboplatin

Lung · NCCP 00568 · version 5

Your treatment journey

Before cycle 1

Blood tests, kidney function, glucose and thyroid testing; folic acid and vitamin B12 support for pemetrexed.

Between cycles

Watch for immune symptoms and chemotherapy effects; they can occur at any time.

Maintenance

Pembrolizumab + pemetrexed every 21 days until progression or unacceptable toxicity, according to the treatment plan.

Cycles 1-4 · day 1

Pembrolizumab + pemetrexed + carboplatin every 21 days.

After cycle 4

Carboplatin stops if the plan is followed and disease/toxicity allow.

Monitoring and support

- Take folic acid exactly as prescribed while on pemetrexed and for the required period after it stops.
- Vitamin B12 is given before pemetrexed and then at protocol-defined intervals.
- Blood count, kidney/liver profile and glucose are checked before each cycle; thyroid tests are repeated during treatment.
- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

05 / when to call Do not wait for the next appointment Use your oncology SOS instructions Call oncology urgently for fever, new breathlessness, persistent diarrhoea, jaundice, severe rash, marked weakness, confusion or any rapidly worsening symptom. Severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Scan to open your treatment portal

Regimen information only; this code contains no patient details.

<https://sactcheck.com/docs/patient/00568/>



Official NCCP source · SACTCheck v0.80.2 · Developed by Dr Paul O'Brien. Patient information prototype for clinical review; follow your oncology team's advice and local emergency instructions.