

Adjuvant abemaciclib + endocrine therapy

Breast · NCCP 00619 · version 4a

Your medicines

Abemaciclib · Endocrine therapy

Your treatment schedule


Continuous tablets alongside endocrine therapy; follow your prescribed plan.

A picture of side effects

Tell your team about new or worsening symptoms. The same symptom can have more than one cause. Numbers connect each symptom box to its location.

Abemaciclib · targeted tablets

Endocrine therapy · additional effects



This map shows selected effects to discuss; it does not show their likelihood or every possible side effect. Blood-count and whole-body effects are marked schematically.

Selected symptoms to help discussion. Tell your team about any new or worsening symptom. See page 2 for the schedule, support and QR.

Adjuvant abemaciclib + endocrine therapy

Breast · NCCP 00619 · version 4a

Your treatment journey

Every day

Abemaciclib 150 mg twice daily, with or without food, at approximately the same times each day unless your dose has been changed.

Alongside it

Continue the endocrine treatment prescribed for you.

First 2 months

Blood count, kidney/liver and bone profile are checked every 2 weeks.

Next 2 months

These tests are usually monthly, then as clinically indicated.

Treatment duration

Abemaciclib is taken continuously for 2 years or until progression/unacceptable toxicity, according to the NCCP source and your treatment plan.

Monitoring and support

- Start anti-diarrhoeal treatment at the first sign of loose stools exactly as advised by your oncology team.
- Swallow tablets whole; do not chew, split or crush.
- Avoid grapefruit/grapefruit juice. If a dose is missed or vomited, take the next dose at the scheduled time rather than an extra dose.
- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

05 / when to call Do not wait for the next appointment Use your oncology SOS instructions Call oncology urgently for fever, persistent or severe diarrhoea, dehydration, new breathlessness, one-sided leg swelling, jaundice, significant bleeding or marked weakness. Severe chest pain or breathing difficulty: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Scan to open your treatment portal

Regimen information only; this code contains no patient details.

<https://sactcheck.com/docs/patient/00619/>



[Official NCCP source](#) · SACTCheck v0.80.2 · Developed by Dr Paul O'Brien. Patient information prototype for clinical review; follow your oncology team's advice and local emergency instructions.