NCCP 00278 · clinician draft

Dose-dense AC → paclitaxel every 2 weeks

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Taxane / combination phase

Active medicines: Paclitaxel

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- Fortnightly paclitaxel: dexamethasone 20 mg orally at 12 and 6 hours before treatment, OR the source IV option 30 minutes before; chlorphenamine 10 mg IV and famotidine 20 mg IV before treatment. The team selects the route / timing; do not substitute the weekly 8 mg first-dose schedule.
- Growth-factor support is required for the dose-dense taxane phase. Consultant-directed steroid reduction / omission must account for reactions and interactions; if aprepitant is added, the source discusses 12 mg treatment-day dexamethasone.
- Dose-dense treatment requires the prescribed G-CSF plan. NCCP requires G-CSF with all cycles and refers to local policy; specify the chosen product, dose and days in the individual prescription.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00278 v7 · p6 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-23 · SACTCheck v0.84.17 · Regimen-level drafts; prescriber and oncology-pharmacy verification required