NCCP 00423 · VERSION 5

AT 50/75

Doxorubicin followed by docetaxel on Day 1 every 21 days, for up to six cycles unless progression or unacceptable toxicity changes the plan.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Heart

Doxorubicin

Baseline assessment; report chest pain, breathlessness or swelling.

3 Hair / scalp


Chemotherapy phase

Hair loss is common. Ask about scalp cooling before treatment where suitable.

5 Nerves / nails

Docetaxel

Report numbness, tingling or painful nails before the next dose.



2 Blood / infection

Chemotherapy phase

Fever, shivers or suddenly unwell: call immediately.

4 Mouth

Chemotherapy phase

Painful ulcers or difficulty drinking need advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects are highlighted. The map is not a complete list; persistent or new symptoms need review even after a medicine stops.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

NCCP 00423 · PRACTICAL SUPPORT

Your treatment, at home.

Important medicine-specific effects

Doxorubicin

- Low counts, infection and mouth ulcers
- Nausea, hair loss and fatigue
- Heart damage, sometimes delayed
- Severe tissue injury if the infusion leaks
- Rare later blood cancers

Docetaxel

- Low blood counts, infection and sepsis
- Hair loss, mouth soreness and fatigue
- Numbness / tingling; sometimes lasting
- Fluid retention and nail changes
- Allergic infusion reaction

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Steroid tablets start before the visit

Follow the docetaxel steroid prescription from the day before treatment. If doses are missed, call the unit before attending; do not improvise a replacement.

A clear anti-sickness plan

Take the prescribed prevention medicines and keep the rescue instructions. Contact the team if nausea or vomiting prevents drinking.

Infection and growth-factor plan

The team decides whether growth-factor injections are needed and gives the exact days. Fever or suddenly feeling unwell needs immediate advice.

Hydration and urinary symptoms

Report new swelling, breathlessness or rapid weight gain. Do not change prescribed steroids yourself.

Before treatment

Discuss fertility, contraception, other medicines and line access. Tell staff immediately about burning or pain at the infusion site.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00423/

Sources and review status are available online. Follow your own prescription and hospital instructions.