Trifluridine/tipiracil (Lonsurf) monotherapy

Oral chemotherapy to help control previously treated colorectal cancer that has spread. This plan is Lonsurf alone, without bevacizumab.

Previously treated metastatic colorectal cancer · NCCP 00382 v4 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

This guide covers Lonsurf on its own for previously treated metastatic colorectal cancer under NCCP 00382a. It does not cover gastric cancer or Lonsurf with bevacizumab. Confirm the exact regimen on your prescription.

My overall treatment

Take prescribed morning and evening doses on Days 1–5 and 8–12, within one hour after meals. No Lonsurf on Days 6–7 or 13–28. The team authorises each new 28-day cycle.

28-day cycle
Two five-day tablet blocks, then recovery. No fixed total number of cycles.

Your treatment schedule

Days 1–5	Days 6–7	Days 8–12	Days 13–28
Prescribed morning/ evening tablets after meals.	No Lonsurf.	Second morning/evening tablet block.	No Lonsurf; recovery, tests and review.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline bloods / urine** Required by regimen	Blood count, kidney and liver tests; urine-protein dipstick.
Each cycle	**Bloods / urine / symptoms** Required by regimen	Repeat before every cycle or as indicated; review recovery and the prescription.
If needed	**Extra blood counts** Individual risk	During a cycle as arranged, especially with kidney impairment or symptoms.
After / review	**Recovery / scans / markers** Individual plan	Team-agreed plan and follow-up of unresolved effects; no fixed scan interval.

Between treatments

- Rest days are not guaranteed safe days: report fever, sudden illness, bleeding or poor intake promptly.
- Do not replace missed, held or vomited doses, double doses or restart without the team’s instructions.

Confirm exact morning/evening tablet quantities and strengths, the calendar, anti-sickness medicines, tests and urgent contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Energy / wellbeing


Trifluridine/tipiracil

Fatigue, dizziness or sudden illness: report changes.

2 Mouth

Trifluridine/tipiracil

Soreness or ulcers can affect eating and drinking.



3 Blood counts

Trifluridine/tipiracil

Infection, anaemia or bleeding may occur; blood tests and urgent symptom reporting matter.

4 Sickness / appetite

Trifluridine/tipiracil

Nausea, vomiting or poor intake can cause dehydration.

5 Kidneys / fluids

Trifluridine/tipiracil

Reduced urine or dehydration needs review; kidney impairment can increase drug toxicity.

6 Bowel

Trifluridine/tipiracil

Diarrhoea, cramps or severe tummy pain: call early.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call your oncology team immediately for a temperature of 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell. Infection can occur without a fever. Call urgently for bleeding, severe diarrhoea, repeated vomiting, inability to drink, very little urine or new breathlessness. Severe breathing difficulty, heavy bleeding or collapse needs 112 / 999.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Previously treated metastatic colorectal cancer · NCCP 00382 v4 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Lonsurf (trifluridine/tipiracil)

28 days · tablets twice daily Days 1–5 and 8–12 · remaining days tablet-free.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.